

# Raising Guinea Fowl

Jacquie Jacob and Tony Pescatore, Animal and Food Sciences

Guinea fowl, native to Africa, were brought to the United States by early settlers. A guinea fowl is definitely not a chicken. It is more active, ranges farther, and flies higher than a chicken. There are three species of guinea fowl:

- **Vulturine** guinea fowl (*Acrylium vulturinum*) are from the semi-arid regions of east Africa. The vulturine is not commonly raised in the United States as it is more sensitive to cold and lacks the hardiness of the common guinea fowl. It has a helmetless head and resembles a vulture.
- **Crested** guinea fowl (*Guttera pucherani*) are from South Africa. They can be distinguished from other guinea fowl by their black head plumes.
- **Helmeted** guinea fowl (*Numididae meleagris*) are the most common species of guinea fowl. They are named for the bony “helmet” on their head (see Figure 1). Only three colors are recognized for exhibition purposes: Pearl, White and Lavender.



Figure 1. A pearl grey helmeted guinea hen. Jacquie Jacob

The remainder of this publication refers to raising the helmeted guinea fowl.

## Why Keep Guinea Fowl?

Guinea fowl are rough, vigorous, hardy, and mostly disease-free game birds. They are increasing in popularity for a variety of reasons.

- **For show:** Many poultry shows include a class of guinea. Guinea fowl can typically be entered in one of two classes—breeding (judged according to a Standard of Perfection) and market (judged according to body conformation as it relates to meat production).
- **For income:** The meat of a young guinea is tender, resembling that of wild game. Guinea fowl meat has many nutritional qualities that make it a valuable addition to the diet. It has 134 calories per 100 grams, which is second only to turkeys, which have 109 calories in the same quantity of meat. Guinea fowl meat is leaner and drier than chicken meat. Guinea fowl are ready to eat at 14 to 16 weeks of age.

Guinea eggs can be eaten just as chicken eggs and should be collected daily if not used for hatching purposes. During the laying season an egg a day is common.

The feathers of guinea fowl are also often sought for use in arts and crafts in such items as hats, dream catchers and decorative gourds.

- **Farm yard watch animal:** Guinea fowl will sound an alarm whenever anything unusual occurs on the farm. Some find this noise a nuisance, but for others the guinea is an effective tool for protecting the farm livestock. Guineas have been proven as watchdogs in non-farm locations, such as junk yards as well. When dogs are used as guard animals, lawsuits are a possibility if an intruder is injured. In a junkyard with trees, guineas roost in the trees at night and will make a ruckus if disturbed, alerting humans of intruders. Losses to theft can be reduced with no law suits. Guinea fowl will also alert other poultry on the farm of danger. The warning call of guineas notifies others when raptors are in the area, giving them time to dash under shelter.
- **Insect control:** The main food for wild guineas is insects. Guineas will consume large amounts of insects, typically leaving the vegetable and flower gardens alone. Unlike chickens, guineas do not scratch the dirt much and thus do very little damage to gardens. Once established on a farm, guineas are able to pretty much fend for themselves, consuming insects, seeds and grasses. They have been used to control deer ticks, wood ticks, grasshoppers, flies, crickets and other insects. They will also eat slugs. Guineas are

used in some areas to reduce the threat of Lyme disease, which is carried by the deer tick population. Guinea fowl will also eat honey bees, so you have to be careful if you also have apiaries. Guinea fowl have been known to stand by a hive and eat the bees as they come out.

- **Rodent control:** The loud call of the guineas has been shown to discourage rodents from invading the area. Flocks of guineas will kill and eat mice and small rats.
- **Snake control:** Some snakes will eat eggs and baby chicks. In areas where snakes are a problem, groups of guineas have even been known to locate and kill snakes before they can cause harm.
- **Unique ornamental value:** Guineas are curious and interesting animals. Some people like to keep them around simply for the entertainment value. Guinea fowl have been described as darting around the yard like little roadrunners. Guinea droppings (guano) are less offensive than most poultry species and are only messy when the birds are sick. The guano is basically dry and washes away easily in the rain, so you do not have to worry about stepping in a pile of droppings when walking around the yard.

## Varieties

There are three principal varieties of domesticated guinea fowl recognized by the American Poultry Associations in their Standard of Perfection: pearl, white, and lavender. In all three varieties, the head and neck are bare but there may be wattles present. The wattles of the male guinea are much larger than those of the female.

- The **pearl variety** is the most popular and typically the one that people recognize. The birds have purplish-gray plumage with dots of white spots giving the look of pearls. Feathers from the pearl variety are often used for ornamental purposes.
- The next most common variety is the **white guinea** (also called African white). As the name indicates, the white guinea has pure white feathers. In addition, its skin is lighter than the other two varieties. These birds are not albino and are the only solid white bird that hatches solid white and not yellow.
- **Lavender guineas** are similar to the pearl, but their plumage is light gray or lavender dotted with white spots.

## Before You Start

Before you start raising any kind of poultry you should check zoning regulations for your area to make sure that you are permitted to raise poultry. This is particularly important when raising guinea fowl because guineas will range and cross the boundaries of a small lot. It is common for urbanites to move into areas zoned agriculture but then complain if someone's animals cross onto their property. Guinea fowl are usually quiet, but can be very noisy if disturbed.

## Care and Management

### Housing

Guinea fowl are often left to fend for themselves, but it is best if they are provided with a shelter to protect them from high winds, rain, cold, sun, and predators. The shelter can be from a facility you build specifically for them or can simply be room allotted to them in a barn that you already have. If the barn is unheated, it is best

## Judging Guinea Fowl

The correct age and gender classifications for guinea fowl in poultry shows are cock, hen, cockerel, and pullet. Guinea fowl are rarely handled by the judge until the final placings are decided upon, and then only those birds which he/she expects to place. Those birds are then examined for any defects or disqualifications not apparent from the aisle. Judges should keep in mind that a female guinea fowl in heavy production may have faded leg color. Duck foot can occur in guinea fowl and judges are advised to be on the lookout for it and to disqualify any bird with a duck foot.

if you do not insulate the house. Insulation tends to keep moisture in more than it keeps the cold out. Allowing moisture to build up in a poultry house can cause respiratory problems.

The floor of the house should be covered with an absorbent bedding material such as hay, straw or wood shavings. If the litter is kept dry, it need not be replaced for several months. Guineas like to roost, so it is important to provide perches.

Do not confine male guineas with chickens if there are roosters in the same flock. However, if your flock is allowed to free range during the day and is only locked up at night, it is okay to keep the guineas and roosters in the same barn. It is also okay to house them together in a short-term emergency, such as a blizzard or other bad weather. If the male guineas are housed with the roosters full time, however, the guineas will chase the roosters, keeping them from food and water.

If you are keeping guineas for egg production (for hatching or human consumption) there should be nest boxes available. Nest boxes designed for chickens will usually work. It is best if you keep your guinea hens in the henhouse until noon each day so that they will lay eggs inside rather than in a hidden nest outside.

Fresh, clean water must be available at all times. A variety of different waterers are available, and whatever you choose should be cleaned daily.

### **Brooding and rearing**

It is best to start with day-old keets (baby guinea fowl). You can purchase them from a local breeder, a local feed mill, or online with delivery through the mail. The main problem with mail orders is that there is often a minimum order required and you may not want to purchase that many keets. The minimum number that can be ordered is usually based on the best survival rate with shipment. Keets coming through the mail are dependent on their combined body heat in order to survive without a heat lamp during shipment. The minimum number is also set to prevent the keets from being tossed around in the shipping box.

Newly hatched birds, including keets, absorb what is left of the yolk that provided food for the developing embryo during incubation. The absorbed yolk provides food to the newly hatched keets, making it unnecessary for them to eat or drink for 48 hours after they are hatched.

Being native to dry areas of Africa, guineas are very susceptible to dampness during their first two weeks and can die from following the mother through dewy grass. After two weeks of age, however, they are probably the most hardy of all domestic land fowl.

Keets may be raised in the same kind of brooder houses and brooders as chicks or poults (baby turkeys). Hovers should be started at 95°F (35°C) for the first two weeks, and then lowered by 5°F (2.75°C) per week thereafter.

After they are fully feathered, guineas are typically able to tolerate extremes in weather fairly well.

### **Sexing**

One of the most often asked questions about poultry is how to tell the males from the females. It is very hard to sex young guineas since they look exactly the same. You can only really sex guineas once they are older. You can tell them apart by:

- **The sounds they make.** The hen makes a two-syllable noise that sounds like she is saying “buckwheat, buckwheat,” “put-rock, put-rock” or “qua-track, qua-track.” This is the only sound that the hen makes that the rooster does not. Most young birds start making these sounds at 6 to 8 weeks of age, but some females do not start calling until much later.
- **The size of the helmet and wattles.** The helmet and wattles of the male are much larger than those of the female.

### **Managing Adults**

If you are purchasing guineas for tick and insect control then you are better off purchasing adult guineas as they require little care and do very well on their own. Clean water should be made available. Guineas enjoy a little scratch feed scattered on the ground. Guineas allowed to roam freely will eat very little feed during the summer months. If you want them for pest control, restricting their feed will encourage them to spend more time eating insects.

<b>Terminology</b>	
Guinea cockerel	A male guinea less than a year old
Guinea cock	An adult male guinea a year or older
Guinea fowl	Proper name for the species
Guinea hen	An adult female guinea a year or older
Helmet	Protuberance on the head of some guinea fowl
Guinea keet	Offspring of a guinea hen and a guinea cock from hatch through 12 weeks of age
Poultry	Domestic fowl raised for meat or eggs
Guinea pullet	A female guinea less than a year old
Varieties	Differences in color patterns or markings
Young guinea	Any guinea 12 to 52 weeks old

It takes guineas a while to get settled into a new home. If you purchase adult guineas, it is best to keep them confined for a week or two to let them get accustomed to their new home. If you let them out right away they may disappear down the street. They should be confined in a pen where they can see the area where they will be living. After the initial 1 to 2 weeks, let one guinea out. Guineas hate to be alone so the single guinea will not go far and will learn its way around the area. After a few days, let a second guinea out to run with the first. If they stay around it is usually safe to let the rest out.

Guineas are very good runners and prefer this method of getting around, including to escape predators, rather than flying. If the adult guineas are let loose to forage freely, space is usually not a problem. If you are going to confine the guineas for any length of time, they

should be given a minimum indoor floor space of 2 to 3 square feet per guinea. The more room the guineas have, the less likely it is that they will be stressed.

If you want to keep your guineas from wandering from a specific area, they need to be kept in covered pens. Guineaes are able to fly at a very early age and will often fly around the farm. Guineaes are strong fliers and often fly 400 to 500 feet at a time.


### Nutrition

Adult guineas are low maintenance. They will forage for themselves and are able to meet most of their nutrition requirements on their own. They will consume a variety of insects (mosquitoes, ticks, and beetles), grass, weed seeds, slugs, worms, and caterpillars.

Guineas need to consume greens in order to maintain good digestion. They will eat grass, dandelions, weeds, etc. Because they are consuming vegetation, it is important to have grit available. It is also good to have oyster shell available free choice.

If you do not allow your guinea fowl to forage, they can be fed a commercial poultry diet. Guineaes need a higher protein feed than chickens but do quite well on regular poultry diets. Keets need a 24 to 26 percent protein ration for the starter feed. Use an unmedicated feed. The protein level should be reduced to 18 to 20 percent for the fifth to eighth weeks. After that, feed a regular laying mash, which usually contains 16 percent protein. If your local feed mill does not sell the feeds in the protein levels you need, you can mix a higher protein diet with a laying mash to get the proper protein level. Guineaes should be fed only mash or crumbles. Pelleted feed is not recommended for feeding guineas.

**Figure 2.** Parts of a guinea fowl



### Body Parts

Body part	Description
<b>Helmet</b>	Pale, waxy appendage protruding from the top of the head. Hard and curves away from the body. In young keets: blackish color; in adults, brown. Also called the topknot, comb, or casque.
<b>Wattles</b>	Fleshy coral red appendages, flat, stiff, and free from wrinkles, hang from either side of the head. The lower edge of the wattles curves from the beak, is much larger and has a cupped appearance.
<b>Head</b>	Wedge-shaped and short, tapering toward the beak. The head is covered with white skin and has a line of very fine hair-like feathers down the back of the head, which stick out and grow upward.
<b>Eyes</b>	Alert, large, round and dark brown.
<b>Neck</b>	Rounded with down-like feathers.
<b>Shanks</b>	Bright orange in keets; changes to mottled orange with brown as they get older. In older adults, they are black.
<b>Toes</b>	Strong and straight
<b>Beak</b>	Short, strong and curved
<b>Wings</b>	Large and carried horizontally
<b>Back</b>	Broad, curves from the neck descending gradually to the tail
<b>Body</b>	Carried low to the ground and rather short

From the Standard of Perfection for Guinea fowl at <http://www.guineafowl.com/standards.html>.

Supplemental greens such as leafy alfalfa should also be given for the guineas to peck at. They will primarily eat the leaves. Remove any leftovers daily to prevent creating a mold problem.

If the guineas are able to get plenty to eat in the area in which the forage, they will not eat much supplemental feed. They do like the occasional treat of wheat, sorghum (milo) or millet. They do not care much for the larger grains and will ignore whole corn kernels. Whole or cracked grains can be used as supplemental grains.

## Health

Guineas are subject to the same diseases and parasites as other poultry and respond to the same treatments.

**Do not** feed medicated poultry diets to guinea fowl. Coccidiostats, such as coban, are toxic to guinea fowls.

## Marketing

The normal marketing season is late summer and throughout the fall. The demand is typically for young birds weighing 1.75 to 2.5 lb (~1 kg) live weight. Many hotels and restaurants in large cities serve guineas at banquet and club dinners as a special delicacy or as a substitute for other game birds.

Guineas are killed and dressed in the same way as chickens. The bones of guineas are relatively small, and the carcass produces a relatively large amount of meat.

## Reproduction

### Selecting Breeders

When selecting breeders from your flock, size and uniform color are important. Both male and female adult guineas weigh 3.0 to 3.5 lb (~1.5 kg). As previously indicated, guinea fowl are sexed by the cry they make and the size of the helmet and wattles (which are larger in the male).

Wild guineas typically mate in pairs, but under domestic conditions, you usually need only one male for each four to five females.

Hens can lay more than 100 eggs per year. Guinea eggs are smaller than chicken eggs; about 1.4 oz (40 g) versus 2 oz (60 g) for a chicken egg.

### Hatching Egg Production

When allowed to incubate eggs naturally, guinea hens typically do not go broody until the nest has around thirty eggs. A good guinea hen will lay an egg a day. If the eggs are removed, she will most likely make a nest elsewhere. If all but four to six eggs are removed from a nest for safekeeping, she may return to the same spot and continue laying. Using this procedure will allow you to collect eggs daily without having to go on a scavenger hunt. Eggs from confined guineas, however, can typically be collected daily with no problems.

## Incubation

Guineas often lay their eggs out in the fields and hatch their young by themselves. If you do find the eggs and wish to incubate them, the eggs should be treated like chicken eggs. There is one big difference, however: Guinea eggs are smaller and have a thicker shell than chicken eggs, making them nearly impossible to candle for fertility until they have been incubated for at least 10 days.

The incubation period for guinea eggs is 26 to 28 days. The eggs can be incubated in most commercially available incubators. If the incubator does not come with guinea-specific instructions, follow the guidelines for pheasants or turkeys.

If using natural incubation, guinea hens typically begin to lay in April or May but do not become broody until later in the season. If you want to hatch the first eggs laid, you can use a broody chicken hen. Usually a bantam hen can incubate 10 to 15 eggs, and a standard size hen can incubate 25 to 30.