

2014 Alfalfa Grazing Tolerance Report

G.L. Olson, S.R. Smith, and G.D. Lacefield, Plant and Soil Sciences, and J.D. Clark, Animal and Food Sciences

Introduction

Alfalfa (*Medicago sativa*) is the highest-yielding, highest-quality forage legume grown in Kentucky. It forms the basis of Kentucky's cash hay enterprise and is an important component in dairy, horse, beef, and sheep diets. Recent emphasis on its use as a grazing crop and the release of grazing-tolerant varieties have raised the following question: Do varieties differ in tolerance to grazing? We have chosen to use the standard tolerance test recommended by the North American Alfalfa Improvement Conference. This test uses continuous heavy grazing to sort out differences in grazing tolerance in a relatively short period of time.

This report summarizes research on the grazing tolerance of alfalfa varieties when subjected to continuous heavy grazing pressure during the grazing season. Table 7 shows a summary of all alfalfa varieties tested in Kentucky during the last 20 years. The UK Forage Extension Web site, at www.uky.edu/Ag/Forage, contains electronic versions of all forage variety testing reports from Kentucky and surrounding states and from a large number of other forage publications.

Important Selection Considerations

Local Adaptation and Seasonal Yield. The variety should be adapted to Kentucky as indicated by good winter survival and good performance across years and locations in replicated yield and grazing trials, such as those presented in this publication. Choose high-yielding, persistent

varieties and varieties that are productive during the desired season of use. Refer to the 2014 Alfalfa Report (or previous years if needed) for yield data on specific varieties of interest.

Seed Quality. Buy premium-quality seed that is high in germination, high in purity, and free from weed seed. Buy certified seed or proprietary seed of an improved variety. An improved variety is one that has performed well in independent trials. Other information on the label will include the test date (which must be within the previous nine months), the level of germination, and percentage of other crop and weed seed. Order seed well in advance of planting time to assure that it will be available when needed.

Description of the Tests

Alfalfa variety tests for grazing tolerance were established in Lexington in the fall of 2010, 2011, 2012 and 2013. The soils at this location are well-drained silt loams and are well-suited to alfalfa. Plots were 5 feet by 20 feet in a randomized complete block design, with each variety replicated six times. In each test,

20 pounds per acre of seed were planted into a prepared seedbed using a disk drill. All seed lots were treated with metalaxyl fungicide and inoculated if not supplied with these treatments. Plots were grazed continuously beginning the first spring after seeding. Grazing pressure was maintained to keep plant height to less than 3 inches. In general, plots were grazed from April until mid-September. Supplemental hay was fed during periods of slowest growth. Visual ratings of percent stand were made in the fall several weeks after the cattle were removed to check stand survival after the grazing season. Ratings were made in the spring prior to grazing to check on winter survival and spring growth. Since trials were seeded in rows, persistence ratings were based on density within a row and not total ground cover. Pests (weeds and insects) were controlled so they would not limit yield or persistence. Fertilizers (lime, P, K, and boron) were applied based on soil test recommendations. In each trial, Alfagraze was the grazing-tolerant check variety, and either Apollo or 5432 was the grazing-intolerant check variety.

Results and Discussion

Weather data for Lexington for 2011, 2012, 2013 and 2014 are presented in Table 1.

Data on percent stand are presented in Tables 2, 3, 4, and 5. Statistical analyses

Table 1. Temperature and rainfall at Lexington, Kentucky, in 2011, 2012, 2013 and 2014

	2011				2012				2013				2014 ²			
	Temp		Rainfall		Temp		Rainfall		Temp		Rainfall		Temp		Rainfall	
	°F	DEP ¹	IN	DEP	°F	DEP	IN	DEP	°F	DEP	IN	DEP	°F	DEP	IN	DEP
JAN	29	-2	2.10	-0.76	38	+7	4.80	+1.94	38	+7	4.50	+1.64	25	-6	2.28	-.58
FEB	39	+4	6.34	+3.13	40	+5	5.39	+2.18	36	+1	1.78	-1.43	30	-5	5.47	+2.26
MAR	47	+3	4.76	+0.36	56	+12	5.64	+1.24	39	-5	5.47	+1.07	39	-5	3.08	-1.32
APR	58	+3	12.36	+8.48	56	+1	3.26	-0.62	55	0	4.46	+0.58	58	+3	5.27	-1.89
MAY	64	0	6.72	+2.25	69	+5	4.02	-0.45	65	+1	5.23	+0.76	66	+2	5.72	+1.25
JUN	74	+2	2.61	-1.05	73	+1	2.42	-1.24	72	0	7.32	+3.66	75	+3	2.93	-0.73
JUL	80	+4	6.29	1.29	81	+5	2.50	-2.50	72	-4	9.33	+4.33	74	-2	3.18	-1.82
AUG	75	0	2.89	-1.04	75	0	1.68	-2.25	72	-3	3.68	-0.25	76	+1	6.53	+2.60
SEP	66	-2	5.52	+2.32	67	-1	6.40	+3.20	67	-1	2.21	-0.99	69	+1	3.63	+4.43
OCT	55	-2	4.10	+1.53	55	-2	2.00	-0.57	55	-2	7.02	+4.45	57	0	5.55	+2.98
NOV	50	+5	9.53	+6.14	43	-2	1.81	-0.65	41	-4	3.06	-0.33				
DEC	41	+5	5.58	+1.60	42	+6	9.57	+4.94	36	0	4.19	+0.21				
Total			68.80	+24.25			49.49	+4.94			58.25	+13.70			44.14	+6.96

¹ DEP is departure from the long-term average.

² 2014 data is for ten months through October.

were performed on all alfalfa yield data (including experimentals) to determine whether the apparent differences are truly due to variety or just due to chance. Varieties not significantly different from the highest numerical value in a column are marked with one asterisk (*). To determine whether two varieties are truly different, compare the difference between the two varieties to the Least Significant Difference (LSD) at the bottom of the column. If the difference is equal to or greater than the LSD, the varieties are truly different when grown under the conditions at a given location. The Coefficient of Variation (CV), which is a measure of the variability of the data, is included for each column of means. Low variability is desirable, and increased variability within a study results in higher CVs and larger LSDs.

Apollo and 5432 have been used widely in trials as the grazing-intolerant varieties. Therefore, the response of these varieties provides a useful measure of the severity of the grazing pressure applied to the plots. In general, types developed for tolerance to grazing tolerated heavy grazing pressure better than hay types. Table 6 summarizes information about distributors, fall dormancy ratings, disease resistance information and persistence across years for all varieties included in these tests.

Table 7 is a summary of stand persistence data from 1994 to 2014 of commercial varieties that have been entered in the Kentucky trials. The data for each specific trial are listed as a percentage of the grazing-tolerant variety Alfagraze. In other words, in each trial Alfagraze is 100 percent—varieties with percentages over

Table 2. Stand persistence of alfalfa varieties sown September 1, 2010, in a cattle grazing tolerance study at Lexington, Kentucky

Variety	Percent Stand								
	2010	2011		2012		2013		2014	
	Oct 14	Mar 15	Nov 7	Mar 23	Oct 29	Mar 28	Oct 17	Apr 3	Oct 9
Commercial Varieties—Available for Farm Use									
TS 4010/A4535	100	100	43	46	44	36	28	17	16*
Alfagraze	99	99	44	31	28	26	15	10	11*
Ameristand 403T	100	99	45	40	35	31	21	10	10*
TS 4007	99	98	39	29	23	20	13	8	9
PGI 424	97	96	37	34	28	23	11	5	5
Apollo	99	99	37	23	19	14	5	4	3
Mean	99	99	41	34	30	25	15	9	9
CV,%	1	2	26	32	39	44	65	58	61
LSD,0.05	1	2	13	13	14	13	12	6	6

*Not significantly different from the highest numerical value in the column, based on the 0.05 LSD.

Table 3. Seedling vigor and stand persistence of alfalfa varieties sown September 13, 2011, in a cattle grazing tolerance study at Lexington, Kentucky

Variety	Seedling Vigor ¹ Oct 11, 2011	Percent Stand						
		2011	2012 ²		2013		2014	
		Oct 11	Mar 23	Oct 10	Mar 21	Oct 15	Apr 3	Oct 9
Commercial Varieties—Available for Farm Use								
Alfagraze 300 RR	4.0	100	97	99	99	73	68	65*
TS 4010/A4535	4.6	100	97	99	99	68	53	65*
Archer III	4.8	100	98	99	99	65	58	63*
Ameristand 403T Plus	3.8	100	100	100	100	66	55	62*
LegenDairy 5.0	4.6	100	96	99	99	63	53	58*
PGI 459	4.5	100	98	98	99	60	51	55*
Ameristand 407TQ	4.4	100	97	99	98	55	45	53*
Alfagraze	3.8	100	99	100	100	71	55	52
Apollo	4.0	100	96	85	99	56	32	37
Experimental Varieties								
TS 4013	4.3	100	98	100	100	73	66	63*
Mena	4.3	100	97	98	99	65	54	57
CV,%	11.2	0	4	12	1	13	19	18
LSD,0.05	0.6	0	5	14	1	10	12	12

¹ Vigor score based on a scale of 1 to 5 with 5 being the most vigorous seedling growth.

² Due to sclerotinia outbreak after sowing this trial and new seedling growth in the spring of 2012, this trial was grazed rotationally during the summer of 2012 to allow establishment of the alfalfa

*Not significantly different from the highest numerical value in the column, based on the 0.05 LSD.

Table 4. Seedling vigor and stand persistence of alfalfa varieties sown August 30, 2012, in a cattle grazing tolerance study at Lexington, Kentucky

Variety	Seedling Vigor ¹ Oct 8, 2012	Percent Stand				
		2012	2013		2014	
		Oct 8	Mar 21	Sep 19	Apr 3	Oct 9
Commercial Varieties—Available for Farm Use						
Alfagraze	3.9	98	99	57	58	56*
Bulldog-505	5.0	100	100	41	38	50*
Ameristand 403T	4.0	99	99	44	44	44*
Apollo	4.7	99	99	47	47	44*
Experimental Varieties						
GA-ALFG-1	4.7	100	100	44	43	42*
Mean	4.5	99	99	47	46	47
CV,%	13.7	1	1	36	33	31
LSD,0.05	0.7	2	1	20	18	18

¹ Vigor score based on a scale of 1 to 5 with 5 being the most vigorous seedling growth.

*Not significantly different from the highest numerical value in the column, based on the 0.05 LSD.

Table 5. Seedling vigor and stand persistence of alfalfa varieties sown September 6, 2013, in a cattle grazing tolerance study at Lexington, Kentucky

Variety	Seedling Vigor ¹ Oct 14, 2013	Percent Stand		
		2013	2014	
		Oct 14	Apr 2	Oct 9
Commercial Varieties—Available for Farm Use				
Bulldog-505	4.4	99	100	100*
Apollo	4.1	98	98	98*
Ameristand 403T	3.9	99	98	98*
Alfagraze	3.6	95	96	96
Experimental Varieties				
GA-ALFG-1	4.8	100	100	100*
DSD08-SC	4.7	99	99	99*
Mean	4.2	98	98	99
CV,%	13.1	2	2	2
LSD,0.05	0.7	2	2	3

¹ Vigor score based on a scale of 1 to 5 with 5 being the most vigorous seedling growth.

*Not significantly different from the highest numerical value in the column, based on the 0.05 LSD.

Table 7. Summary of Kentucky alfalfa grazing trials 1994-2014 (stand persistence shown as a percent of the grazing tolerant Alfagrazee)

Variety	Proprietor	Variety Characteristics ¹										Lexington										Mean ⁵ (#trials)				
		Disease Resistance ²					APH					1994 ^{3,4}	1996	1997	1998	2000	2000	2001	2004	2005	2006		2008	2009	2010	2011
		FD	Bw	Fw	An	PRR	APH	3yr ⁶	3yr	4yr	3yr	2yr	3yr	3yr	3yr	4yr	3yr	4yr	4yr	4yr	4yr		4yr	4yr	4yr	3yr
ABT 205	W-L Research	2	HR	HR	HR	HR	R	HR	R	HR	R	94	84												89(2)	
ABT 350	W-L Research	3	HR	HR	HR	HR	HR	HR	HR	HR	HR	46	69												83(5)	
ABT 405	W-L Research	4	HR	HR	HR	HR	R	R	R	R	R	71	129	100	100	100	100	100	100	100	100	100	100	100	100(14)	
Alfagrazee	Americas Alfalfa	2	MR	R	MR	R	-	HR	HR	HR	HR	100	100	100	100	100	100	100	100	100	100	100	100	100	-	
Alfagrazee300 RR	Americas Alfalfa	3	HR	R	HR	HR	HR	HR	HR	HR	HR	120	53	56	56	85	26	85	125	141	144	50	91	125	78(6)	
Amerigraze 401+Z	Americas Alfalfa	4	HR	HR	HR	HR	R	HR	HR	HR	HR														107(4)	
Ameristand 403T	Americas Alfalfa	4	HR	HR	HR	HR	HR	HR	HR	HR	HR														126(9)	
Ameristand 403TPlus	Americas Alfalfa	4	HR	HR	HR	HR	HR	HR	HR	HR	HR														119	
Ameristand 407TQ	Americas Alfalfa	4	HR	HR	HR	HR	HR	HR	HR	HR	HR														102	
Apollo	Americas Alfalfa	4	R	R	R	R	R	R	R	R	R	48	75	33	47	17	31	25	36	27	25	17	27	71	36(13)	
Arc (certified)	Public	4	LR	MR	HR	HR	-	-	-	-	-	38													-	
Archer III	Americas Alfalfa	5	HR	HR	HR	HR	HR	HR	HR	HR	HR														77(2)	
Baralfa 54	Barenbrug USA	-	R	HR	HR	HR	HR	HR	HR	HR	HR	68		78									33		-	
Cut-n-Graze	Americas Alfalfa	3	HR	HR	HR	HR	HR	HR	HR	HR	HR														-	
FK 421	Donley Seed Co.	4	HR	H	H	H	H	H	H	H	H							100							-	
Feast	Gaist Seeds	3	HR	HR	HR	HR	R	HR	HR	HR	HR	146			87	92									108(3)	
Fortress	Syngenta	3	R	R	R	R	R	R	R	R	R	40	71												56(2)	
Gold Plus	PGI Alfalfa	4	HR	HR	HR	HR	R	HR	HR	HR	HR			81											-	
GrazeKing	FFR/Southern States	5	MR	HR	HR	R	R	R	R	S	R	91	41					50							61(3)	
Haygrazer	Great Plains Research	4	HR	HR	R	R	MR	R	R	MR	R	75	39			38									51(3)	
Integrity	PGI Alfalfa	4	HR	HR	HR	HR	HR	HR	HR	HR	HR							172							-	
Legacy	Green Seed	4	R	R	R	R	R	R	R	R	R	32													-	
Legendairy5.0	Croplan Genetics	3	HR	HR	HR	HR	HR	HR	HR	HR	HR														56(2)	
Magnagrazee	Dairyland Seed Co.	3	HR	HR	HR	R	HR	R	HR	-	HR	56													-	
Pasture Plus	MBS	3	HR	HR	R	HR	HR	MR	HR	MR	HR	60													-	
PGI 424	Producers Choice	4	HR	HR	HR	HR	HR	HR	HR	HR	HR														-	
PGI 459	Producers Choice	4	HR	HR	HR	HR	HR	HR	HR	HR	HR												45	106	62(2)	
Pioneer 98	Pioneer	3	HR	R	HR	R	-	HR	R	-	HR			56											-	
ProGro	MBS Inc.	4	HR	HR	R	HR	HR	MR	HR	MR	HR	71		81											-	
Quantum	ABI Alfalfa	2	HR	HR	HR	HR	R	HR	R	HR	HR														-	
Rebel	Target Seed	4	HR	HR	HR	HR	HR	HR	HR	HR	HR												79		-	
Rugged	Target Seed	3	HR	HR	HR	HR	HR	HR	HR	HR	HR												146		-	
Rushmore	Syngenta	4	HR	HR	HR	HR	HR	HR	HR	HR	HR	32													-	
Saranac AR (cert.)	Public	4	MR	R	HR	LR	-	HR	LR	-	HR	77						100							89(2)	
Spredor 3	Syngenta	1	HR	HR	R	MR	S	MR	S	MR	S	71	123	75					68						96(4)	
Spredor 4	Syngenta	2	HR	HR	HR	HR	R	HR	R	HR	HR											25			-	
Stampede	Allied Seed	3	HR	R	R	HR	R	HR	R	HR	R	73													-	
TS 4007	Producers Choice	4	HR	R	HR	HR	HR	HR	HR	HR	HR														-	
TS 4010/A4535	Producers Choice	4	HR	R	HR	HR	HR	HR	HR	HR	HR														-	
Triple Trust 450	ABI/America's Alfalfa	5	HR	HR	HR	HR	HR	HR	HR	HR	HR	95							145						118(3)	
Wintergreen	ABI Alfalfa	3	HR	HR	HR	HR	R	HR	HR	R	HR		57	72											75(3)	
WL 326GZ	W-L Research	4	HR	HR	HR	HR	HR	HR	HR	HR	HR	118		88											103(2)	
115 Brand	Monsanto	3	HR	HR	R	HR	R	HR	R	HR	R					56	85								71(2)	
5373	Pioneer	4	HR	HR	HRT	MR	LR	MR	LR	MR	LR	21													-	
5432	Pioneer	4	HR	HR	-	MR	-	MR	-	MR	-							51							-	

¹ Variety characteristics: FD = fall dormancy, Bw = bacterial wilt, Fw = fusarium wilt, An = anthracnose, PRR = phytophthora root rot, APH = aphanomyces root rot. Information provided by seed companies.

² Disease resistance: S = susceptible, LR = low resistance, MR = moderate resistance, R = resistance, HR = high resistance.

³ Year trial was established.

⁴ Use this summary table as a guide in making variety decisions, but refer to specific yearly reports to determine statistical differences in stand persistence between varieties. To find actual persistence ratings, look in the yearly report for the final year of each specific test. For example, the Lexington trial planted in 1996 was grazed for 3 years so final persistence report would be "1999 Alfalfa Grazing Tolerance Report" archived in the KY Forage website at <www.uky.edu/Ag/Forage>.

⁵ Mean only presented when respective variety was included in two or more trials.

⁶ Number of years of data.



Mention or display of a trademark, proprietary product, or firm in text or figures does not constitute an endorsement and does not imply approval to the exclusion of other suitable products or firms.
 The College of Agriculture, Food and Environment is an Equal Opportunity Organization.
 12-2014