

# 2014 Summer Annual Grass Report

G.L. Olson, S.R. Smith, and G.D. Lacefield, Plant and Soil Sciences

## Introduction

Summer annual grasses provide an important forage crop option for producers in Kentucky. These grasses are mainly used as emergency or supplemental hay and pasture crops, but little information is available on their yield potential. The purpose of this publication is to summarize the University of Kentucky 2011-2014 forage yield trials with sudangrass, sorghum/sudangrass, millets, and teff.

Sudangrass (*Sorghum bicolor* ssp. *drummondii*) is a rapidly growing annual grass in the sorghum family. It is medium yielding and well suited for grazing or hay because of its smaller stem size. Sudangrass regrows quickly after harvest and can be grazed several times during summer and early fall.

Sorghum x sudangrass hybrids are more vigorous and slightly higher yielding than sudangrass. A larger stem size makes these hybrids less useful for hay; therefore, they are commonly used for baleage and grazing.

Forage sorghum is used primarily as silage for livestock and is typically a one cut crop. It grows 9-12 feet tall and is harvested when the seed is in the dough stage.

Pearl millet (*Pennisetum glaucum*) is the most widely grown type of millet. It is well adapted to production systems characterized by drought, low soil fertility, and high temperature. It is higher yielding than foxtail millet and regrows rapidly after harvest if an 8- to 10-inch stubble height is left. Dwarf varieties, which are leafier and better suited for grazing, are available.

Teff, also referred to as summer lovegrass (*Eragrostis tef*), is a warm-season annual grass native to Ethiopia and has been used as a grain crop for thousands of years. Recently, there has been considerable interest in teff as a forage crop. It is high quality, palatable, and fine-stemmed and, therefore, makes excellent hay.

## Considerations in Selecting a Summer Annual Variety

The major factor in selecting a variety of summer annual grass is yield, both total and seasonal. Growth after first cutting is strongly dependent on available moisture and nitrogen fertilization. Summer annual grasses generally have different characteristics and uses. Pearl millets vary considerably in height and can be used for both pasture and hay. Pearl millet has the advantage of not producing prussic acid (HCN or cyanide). Sudangrass and sorghum-sudangrass hybrids are related grasses (in the sorghum family) and can produce prussic acid immediately after frost or when immature shoots are grazed during severe drought. Sudangrasses are considered to have the least potential for prussic acid poisoning. Sudangrass has smaller, finer stems than sorghum-sudangrass hybrids, which

have finer stems than forage sorghums. Consequently, sudangrasses are more easily cured for hay. Pearl millets, sudangrass, sorghum-sudangrass, and teff are typically harvested multiple times during the growing season, and foxtail millet is harvested only once. For more detailed management recommendations refer to *Producing Summer Annual Grasses for Emergency or Supplemental Forage* (AGR-88), and *Teff*, which can be found at [www.uky.edu/Ag/Forage](http://www.uky.edu/Ag/Forage) under "Publications" in the "Grass" species.

## Description of the Tests

This report summarizes studies at Lexington (three in 2011, three in 2012, five in 2013 and five in 2014). The soil at Lexington (Maury) is a well-drained silt loam and is well suited to annual grass production. Plots were 5 feet x 20 feet in a randomized complete block design with four replications with a harvested area of 5 feet by 15 feet. All trials were sown into a prepared seedbed using a disk drill at the following rates (lb/acre): sudangrass (25), sorghum-sudangrass (30), forage sorghum (8), pearl millet (20), and teff (5 for uncoated, 8 for coated). Plots were harvested with a sickle-type forage plot harvester. Cutting height was 4 inches

**Table 1. Temperature and rainfall at Lexington, Kentucky in 2011, 2012, 2013 and 2014.**

	2011				2012				2013				2014 <sup>2</sup>			
	Temp		Rainfall		Temp		Rainfall		Temp		Rainfall		Temp		Rainfall	
	°F	DEP <sup>1</sup>	IN	DEP	°F	DEP	IN	DEP	°F	DEP	IN	DEP	°F	DEP	IN	DEP
JAN	29	-2	2.10	-0.76	38	+7	4.80	+1.94	38	+7	4.50	+1.64	25	-6	2.28	-.58
FEB	39	+4	6.34	+3.13	40	+5	5.39	+2.18	36	+1	1.78	-1.43	30	-5	5.47	+2.26
MAR	47	+3	4.76	+0.36	56	+12	5.64	+1.24	39	-5	5.47	+1.07	39	-5	3.08	-1.32
APR	58	+3	12.36	+8.48	56	+1	3.26	-0.62	55	0	4.46	+0.58	58	+3	5.27	-1.89
MAY	64	0	6.72	+2.25	69	+5	4.02	-0.45	65	+1	5.23	+0.76	66	+2	5.72	+1.25
JUN	74	+2	2.61	-1.05	73	+1	2.42	-1.24	72	0	7.32	+3.66	75	+3	2.93	-0.73
JUL	80	+4	6.29	1.29	81	+5	2.50	-2.50	72	-4	9.33	+4.33	74	-2	3.18	-1.82
AUG	75	0	2.89	-1.04	75	0	1.68	-2.25	72	-3	3.68	-0.25	76	+1	6.53	+2.60
SEP	66	-2	5.52	+2.32	67	-1	6.40	+3.20	67	-1	2.21	-0.99	69	+1	3.63	+4.43
OCT	55	-2	4.10	+1.53	55	-2	2.00	-0.57	55	-2	7.02	+4.45	57	0	5.55	+2.98
NOV	50	+5	9.53	+6.14	43	-2	1.81	-0.65	41	-4	3.06	-0.33				
DEC	41	+5	5.58	+1.60	42	+6	9.57	+4.94	36	0	4.19	+0.21				
Total			68.80	+24.25			49.49	+4.94			58.25	+13.70			44.14	+6.96

<sup>1</sup> DEP is departure from the long-term average.

<sup>2</sup> 2014 data is for ten months through October.

for teff and 6 inches for millet, sudangrass and sorghum-sudangrass. The forage sorghum was harvested by hand (5 feet by 5 feet in the center of the plot in 2013 and the center 15 foot row in 2014). Fresh weight samples were taken at each harvest to calculate percent dry matter production. All tests were managed for establishment, fertility, pest control, and harvest according to University of Kentucky Cooperative Extension Service recommendations. Pests were controlled so that they would not limit yield. See individual yield tables for nitrogen application.

## Results and Discussion

Weather data for Lexington is presented in Table 1.

Yield data (on a dry matter basis) for all tests are reported in Tables 3 through 18. Varieties are listed in order from highest to lowest total production. Yields are given by cutting and as a total for the year. Statistical analyses were performed on all yield data to determine if the apparent differences are truly due to variety or just due to chance. Varieties not significantly different from the highest numerical value in a column are marked with one asterisk (\*). To determine if two varieties are truly different, compare the difference between the two varieties to the Least Significant Difference (LSD) at the bottom of the column. If the difference is equal to or greater than the LSD, the varieties are truly different when grown under the conditions at a given location. The Coefficient of Variation (CV), which is a measure of the variability of the data, is included for each column of means. Low variability is desirable, and increased variability within a study results in higher CVs and larger LSDs.

Tables 19, 20, 21, and 22 are summaries of yield data from 2008 to 2014 of commercial varieties that have been entered in the Kentucky trials. The data are listed as a percentage of the mean of the commercial varieties entered in each specific trial. In other words, the mean for each trial is 100 percent—varieties with percentages over 100 yielded better than average, and varieties with percentages less than 100 yielded lower than average. Direct, statistical comparisons of varieties cannot be made using the summary Tables 19, 20, 21, and 22, but these comparisons do help to identify varieties for further consideration. Varieties that have performed better than average over many years and at several locations have very stable performance; others may have performed very well in wet years or on particular soil types.

## Summary

Summer annual grasses can be an important supplemental source of pasture, hay, and silage in Kentucky. Varieties should be selected for their seasonal and total yield characteristics and for their suitability for the method of harvest to be employed (pasture, hay, or silage). Make sure seed of the chosen variety is properly labeled and will be available when needed.

**Table 2. Descriptive scheme for the stages of development in perennial forage grasses.**

Code	Description	Remarks
<b>Leaf development</b>		
11	First leaf unfolded	Applicable to regrowth of established (plants) and to primary growth of seedlings.
12	2 leaves unfolded	Further subdivision by means of leaf development index (see text).
13	3 leaves unfolded	
•	•••••	
19	9 or more leaves unfolded	
<b>Sheath elongation</b>		
20	No elongated sheath	Denotes first phase of new spring growth after overwintering. This character is used instead of tillering which is difficult to record in established stands.
21	1 elongated sheath	
22	2 elongated sheaths	
23	3 elongated sheaths	
•	•••••	
29	9 or more elongated sheaths	
<b>Tillering (alternative to sheath elongation)</b>		
21	Main shoot only	Applicable to primary growth of seedlings or to single tiller transplants.
22	Main shoot and 1 tiller	
23	Main shoot and 2 tillers	
24	Main shoot and 3 tillers	
•	•••••	
29	Main shoot and 9 or more tillers	
<b>Stem elongation</b>		
31	First node palpable	More precisely an accumulation of nodes. Fertile and sterile tillers distinguishable.
32	Second node palpable	
33	Third node palpable	
34	Fourth node palpable	
35	Fifth node palpable	
37	Flag leaf just visible	
39	Flag leaf ligule/collar just visible	
<b>Booting</b>		
45	Boot swollen	
<b>Inflorescence emergence</b>		
50	Upper 1 to 2 cm of inflorescence visible	
52	¼ of inflorescence emerged	
54	½ of inflorescence emerged	
56	¾ of inflorescence emerged	
58	Base of inflorescence just visible	
<b>Anthesis</b>		
60	Preanthesis	Inflorescence-bearing internode is visible. No anthers are visible.
62	Beginning of anthesis	First anthers appear.
64	Maximum anthesis	Maximum pollen shedding.
66	End of anthesis	No more pollen shedding.
<b>Seed ripening</b>		
75	Endosperm milky	Inflorescence green
85	Endosperm soft doughy	No seeds loosening when inflorescence is hit on palm.
87	Endosperm hard doughy	Inflorescence losing chlorophyll; a few seeds loosening when inflorescence hit on palm
91	Endosperm hard	Inflorescence-bearing internode losing chlorophyll; seeds loosening in quantity when inflorescence hit on palm.
93	Endosperm hard and dry	Final stage of seed development; most seeds shed.

Smith, J. Allan, and Virgil W. Hayes. 1981. p. 416-418. 14th International Grasslands Conference Proc. 1981. June 14-24, 1981, Lexington, Kentucky.

**Table 3. Dry matter yields, percent stand, seedling vigor ,maturity and stand height of sudangrass varieties sown May 25, 2011 at Lexington, Kentucky.**

Variety	Proprietor/ Distributor	Seedling Vigor <sup>1</sup> Jun 16	Percent Stand Jun 16	Maturity <sup>2</sup>		Plant Height (inches)				Yield (DM tons/acre)				
				Jun 27	Jul 18	Jun 27	Jul 18	Aug 8	Sep 20	Jun 28	Jul 18	Aug 8	Sep 20	Total
<b>Commercial Varieties—Available for Farm Use</b>														
ProMax BMR	Ampac Seed	4.5	99	2.3	2.5	34	41	44	42	0.53	1.05	1.17	0.80	3.54*
SS130 BMR	Cal/West Seeds	3.8	99	1.5	2.0	27	33	38	29	0.49	1.00	1.02	0.67	3.18*
Monarch V	Public	5.0	100	2.0	1.3	33	32	33	29	0.64	0.94	0.81	0.62	3.01*
Hayking BMR	Cal/West Seeds	3.5	97	1.8	3.0	26	41	40	32	0.38	0.92	1.03	0.67	3.00*
Enorma BMR	Cal/West Seeds	3.3	97	1.3	2.3	25	37	41	32	0.37	0.92	0.96	0.66	2.91
Piper	Public	4.8	100	2.0	1.8	33	34	36	30	0.52	0.96	0.88	0.55	2.90
Mean		4.1	99	1.8	2.1	30	36	39	32	0.49	0.96	0.98	0.66	3.09
CV,%		11.6	1	25.5	37.4	12	10	10	21	20.12	10.17	14.16	22.80	13.19
LSD,0.05		0.7	2	0.7	1.2	5	6	6	10	0.15	0.15	0.21	0.23	0.61

<sup>1</sup> Vigor score based on a scale of 1 to 5 with 5 being the most vigorous seedling growth.

<sup>2</sup> Maturity rating scale: 37 = flag leaf emergence, 45 = boot swollen, 50 = beginning of inflorescence emergence, 58 = complete emergence of inflorescence, 62 = beginning of pollen shed. See Table 2 for complete scale.

\*Not significantly different from the highest numerical value in the column, based on the 0.05 LSD.

Nitrogen application: 30 lb/A on June 2, 60 lb/A on June 28 and 40 lb/A on July 18 of actual nitrogen.

The following is a list of University of Kentucky Cooperative Extension publications related to ryegrass management. They are available from your county Extension office and are listed in the “Publications” section of the UK Forage Web site, [www.uky.edu/Ag/Forage](http://www.uky.edu/Ag/Forage).

- Lime and Fertilizer Recommendations (AGR-1)
- Grain and Forage Crop Guide for Kentucky (AGR-18)
- Establishing Forage Crops (AGR-64)

- Producing Summer Annual Grasses for Emergency or Supplemental Forage (AGR-88)
- Forage Identification and Use Guide (AGR-175)
- Extending Grazing and Reducing Stored Feed Needs (AGR-199)

### About the Authors

G.L. Olson is a research specialist and S.R. Smith and G.D. Lacefield are Extension professors of Forages.

**Table 4. Dry matter yields, seedling vigor, percent stand, maturity and stand height of sudangrass varieties sown May 10, 2012 at Lexington, Kentucky.**

Variety	Proprietor/ Distributor	Seedling Vigor <sup>1</sup> Jun 4	Percent Stand Jun 4	Maturity <sup>2</sup>				Plant Height (inches)				Yield (DM tons/acre)				
				Jun 21	Jul 24	Aug 13	Sep 27	Jun 21	Jul 24	Aug 13	Sep 27	Jun 21	Jul 24	Aug 13	Sep 27	Total
<b>Commercial Varieties—Available for Farm Use</b>																
AS9301 BMR	Alta Seeds/ Ramer Seeds	3.6	96	30.8	45.0	31.3	54.8	34	38	29	41	0.87	0.96	0.94	1.41	4.19*
Piper	Public	5.0	100	32.5	54.0	32.5	46.3	44	43	32	35	0.99	1.00	0.86	0.86	3.71
Hayking BMR	Cal/West Seeds	3.1	98	32.0	57.0	32.3	50.5	40	44	32	38	0.83	0.97	0.86	0.81	3.46
ProMax BMR	Ampac Seed	3.1	98	32.5	59.0	33.3	50.8	41	44	32	37	0.78	0.97	0.81	0.87	3.43
Monarch V	Cal/West Seeds	4.0	100	32.0	47.8	32.3	48.0	40	38	29	32	0.86	0.92	0.71	0.81	3.30
Enorma BMR	Cal/West Seeds	2.8	97	32.0	46.3	32.3	50.3	35	40	30	38	0.73	0.97	0.83	0.74	3.27
<b>Experimental Varieties</b>																
CW5-43-29	Cal/West Seeds	3.0	97	32.0	59.0	32.8	59.5	40	48	36	43	0.79	1.04	0.93	0.99	3.75
CW6-43-50	Cal/West Seeds	3.4	99	32.3	60.5	33.3	53.8	38	47	33	38	0.78	1.00	0.83	0.97	3.59
Mean		3.5	98	32.0	53.6	32.5	51.7	39	43	32	38	0.83	0.98	0.85	0.93	3.59
CV,%		14.0	2	2.4	7.1	2.9	8.8	6	9	9	11	9.03	8.38	9.72	16.27	6.99
LSD,0.05		0.7	3	1.1	5.6	1.4	6.7	4	5	4	6	0.11	0.12	0.12	0.22	0.37

<sup>1</sup> Vigor score based on a scale of 1 to 5 with 5 being the most vigorous seedling growth.

<sup>2</sup> Maturity rating scale: 37 = flag leaf emergence, 45 = boot swollen, 50 = beginning of inflorescence emergence, 58 = complete emergence of inflorescence, 62 = beginning of pollen shed. See Table 2 for complete scale.

\*Not significantly different from the highest numerical value in the column, based on the 0.05 LSD.

Nitrogen application: 60 lb/A on May 11, 50 lb/A on July 26 and 50 lb/A on August 14 of actual nitrogen.

Rainfall deficit: May-August rainfall was 10.62 inches; rainfall deficit during this period in 2012 was -6.44 inches.

**Table 5. Dry matter yields, seedling vigor, percent stand, maturity and stand height of sudangrass varieties sown May 28, 2013 at Lexington, Kentucky.**

Variety	Proprietor/ Distributor	Seedling Vigor <sup>1</sup> Jun 20	Percent Stand Jun 20	Maturity <sup>2</sup>			Height (inches)				Yield (DM tons/acre)				
				Jul 8	Aug 7	Sep 9	Jul 8	Aug 7	Sep 9	Oct 21	Jul 8	Aug 7	Sep 9	Oct 21	Total
<b>Commercial Varieties—Available for Farm Use</b>															
SS130 BMR	Cal/West Seeds	4.6	97	32.0	33.3	42.3	43	49	50	16	1.37	1.74	1.69	0.39	5.19*
Piper	Public	4.8	99	32.3	33.5	33.8	46	58	54	24	1.37	1.69	1.53	0.47	5.07*
ProMax BMR	Ampac Seed	4.4	96	32.3	34.0	36.8	44	60	57	20	1.35	1.73	1.52	0.38	4.98*
Monarch V	Public	5.0	99	32.3	33.3	36.3	44	52	46	17	1.40	1.66	1.37	0.33	4.76*
Hayking BMR	Cal/West Seeds	3.9	95	32.0	33.8	39.5	44	55	52	21	1.18	1.60	1.52	0.36	4.66*
Enorma BMR	Cal/West Seeds	3.5	97	32.0	33.5	42.3	40	53	53	17	1.17	1.50	1.44	0.28	4.39
Mean		4.4	97	32.1	33.5	38.5	43	54	52	19	1.31	1.65	1.51	0.37	4.84
CV,%		11.5	3	1.1	1.8	13.2	8	10	7	9	17.19	12.86	7.64	17.29	9.62
LSD,0.05		0.8	4	0.5	0.9	7.6	5	9	6	3	0.34	0.32	0.17	0.10	0.70

<sup>1</sup> Vigor score based on a scale of 1 to 5 with 5 being the most vigorous seedling growth.

<sup>2</sup> Maturity rating scale: 37 = flag leaf emergence, 45 = boot swollen, 50 = beginning of inflorescence emergence, 58 = complete emergence of inflorescence, 62 = beginning of pollen shed. See Table 2 for complete scale.

\*Not significantly different from the highest numerical value in the column, based on the 0.05 LSD.

Nitrogen application: 50 lb/A of actual nitrogen on July 22 and Aug 8.

**Table 6. Dry matter yields, seedling vigor, stand rating, maturity and plant height of sudangrass varieties sown May 21, 2014 at Lexington, Kentucky.**

Variety	Proprietor/ Distributor	Seedling Vigor <sup>1</sup> Jun 17	Percent Stand Jun 17	Maturity <sup>2</sup>		Plant Height (in)			Yield (DM tons/acre)				
				Jul 8	Aug 11	Jul 8	Aug 11	Sep 12	Jul 8	Aug 11	Sep 12	Oct 29	Total
<b>Commercial Varieties—Available for Farm Use</b>													
Trudan Headless	Chromatin	5.0	100	33.0	32.0	38	28	41	1.57	1.15	1.53	0.46	4.71*
Monarch V	Public	4.3	100	46.3	43.5	42	36	32	1.50	1.30	1.21	0.37	4.38*
SS130 BMR	Cal/West Seeds	4.4	99	39.0	46.3	39	37	35	1.45	1.29	1.16	0.34	4.24*
Promax BMR	Ampac Seed	3.8	97	44.3	48.8	41	41	37	1.22	1.27	1.10	0.41	4.00*
Hayking BMR	Cal/West Seeds	3.0	94	39.0	45.0	38	40	35	1.13	1.23	1.05	0.27	3.69
Piper	Public	3.9	98	42.0	45.0	41	38	32	1.20	1.02	0.91	0.43	3.57
Enorma BMR	Cal/West Seeds	3.3	97	43.0	46.3	39	37	33	1.11	1.09	0.90	0.24	3.33
Mean		3.9	98	40.9	43.8	40	37	35	1.31	1.19	1.12	0.36	3.99
CV,%		15.9	2	8.7	4.6	14	9	6	23.91	9.75	15.56	36.65	13.13
LSD,0.05		0.9	3	5.3	3.0	8	5	3	0.47	0.17	0.26	0.20	0.78

<sup>1</sup> Vigor score based on a scale of 1 to 5 with 5 being the most vigorous seedling growth.

<sup>2</sup> Maturity rating scale: 37 = flag leaf emergence, 45 = boot swollen, 50 = beginning of inflorescence emergence, 58 = complete emergence of inflorescence, 62 = beginning of pollen shed. See Table 2 for complete scale.

\*Not significantly different from the highest numerical value in the column, based on the 0.05 LSD.

Nitrogen application: 50 lb/A of actual nitrogen on May 27, July 11 and August 15.

**Table 7. Dry matter yields, percent stand, seedling vigor, maturity and stand height of sorghum-sudangrass varieties sown May 25, 2011 at Lexington, Kentucky.**

Variety	Proprietor/ Distributor	Seedling Vigor <sup>1</sup> Jun 16	Percent Stand Jun 16	Maturity <sup>2</sup>		Plant Height (inches)				Yield (DM tons/acre)					
				Jun 30	Jul 22	Jun 30	Jul 22	Aug 15	Sep 20	Jun 30	Jul 22	Aug 15	Sep 20	Total	
<b>Commercial Varieties—Available for Farm Use</b>															
Sweet-For-Ever	Gayland Ward Seed	3.5	100	30.0	32.3	30	44	33	29	0.59	1.80	1.12	0.88	4.38*	
SS211	Southern States	3.8	96	31.0	32.8	40	49	44	29	0.68	1.58	1.22	0.67	4.15*	
NutraPlus BMR	Cisco	5.0	100	30.3	31.8	36	35	33	24	0.84	1.44	1.17	0.64	4.10*	
Super Sugar	Gayland Ward Seed	4.3	97	31.0	32.8	40	48	45	29	0.70	1.51	1.17	0.72	4.09*	
Special Effort	Cisco	3.8	77	30.8	32.8	36	44	38	28	0.67	1.42	1.10	0.58	3.77*	
GW300BMR	Gayland Ward Seed	3.3	93	30.3	32.5	34	44	37	25	0.55	1.44	1.01	0.54	3.53	
Mean		3.9	94	30.5	32.5	36	44	39	27	0.67	1.53	1.13	0.67	4.00	
CV,%		10.4	19	1.1	1.9	5	9	6	7	9.56	12.09	14.14	13.81	10.36	
LSD,0.05		0.6	27	0.5	0.9	3	6	3	3	0.10	0.28	0.24	0.14	0.63	

<sup>1</sup> Vigor score based on a scale of 1 to 5 with 5 being the most vigorous seedling growth.

<sup>2</sup> Maturity rating scale: 37 = flag leaf emergence, 45 = boot swollen, 50 = beginning of inflorescence emergence, 58 = complete emergence of inflorescence, 62 = beginning of pollen shed. See Table 2 for complete scale.

\*Not significantly different from the highest numerical value in the column, based on the 0.05 LSD.

Nitrogen application: 30 lb/A on June 2, 60 lb/A on July 1, 40 lb/A on July 27 and Aug 19 of actual nitrogen.

**Table 8. Dry matter yields, seedling vigor, percent stand, maturity and stand height of sorghum-sudangrass and pearl millet (PM) varieties sown May 10, 2012 at Lexington, Kentucky.**

Variety	Proprietor/ Distributor	Seedling Vigor <sup>1</sup> Jun 4	Percent Stand Jun 4	Maturity <sup>2</sup>			Plant Height (inches)			Yield (DM tons/acre)			
				Jun 27	Jul 30	Sep 27	Jun 27	Jul 30	Sep 27	Jun 27	Jul 30	Sep 27	Total
<b>Commercial Varieties—Available for Farm Use</b>													
Vita-Cane	Gayland Ward Seed	4.8	100	31.3	44.0	62.0	38	38	44	0.90	1.20	1.52	3.61*
Super Sugar	Gayland Ward Seed	4.6	98	31.0	44.5	62.0	38	38	53	0.77	1.17	1.55	3.49*
Special Effort	Cisco	4.4	96	31.0	50.5	60.0	37	39	44	0.80	1.08	1.56	3.44*
SS220 BMR	Southern States	3.0	64	31.0	37.8	54.0	32	48	47	0.48	1.28	1.59	3.35*
Sweet-For-Ever	Gayland Ward Seed	4.6	99	31.0	31.5	46.0	28	39	40	0.63	1.21	1.38	3.21*
NutraPlus BMR	Cisco	4.6	93	30.5	50.8	56.0	34	44	41	0.69	1.32	1.16	3.17*
SS211	Southern States	2.9	53	31.0	43.5	47.8	29	50	44	0.39	1.08	1.32	2.79
AS6402 BMR	Alta Seeds/Ramer Seed	3.8	75	30.0	32.0	53.0	26	38	34	0.46	1.10	1.17	2.73
Pennleaf Hybrid (PM)	Pennington Seed	2.0	99	29.0	38.8	75.0	17	24	27	0.36	0.96	1.28	2.59
GW 2120	Gayland Ward Seed	4.0	97	31.0	36.8	46.8	27	33	28	0.64	1.12	0.73	2.48
Sweet-For-Ever BMR	Gayland Ward Seed	3.4	93	30.5	32.3	42.0	27	41	29	0.50	0.96	0.87	2.33
GW 300 BMR	Gayland Ward Seed	3.9	91	30.5	42.0	47.3	30	46	34	0.49	1.10	0.74	2.33
PP102M Hybrid (PM)	Cisco	2.0	97	29.5	60.0	75.0	19	41	37	0.35	0.88	0.83	2.05
Tifleaf III Hybrid (PM)	Gayland Ward Seed	2.4	100	29.5	58.5	75.0	19	37	37	0.31	0.87	0.80	1.98
Mean		3.6	89	30.5	43.1	57.5	28	40	38	0.55	1.10	1.18	2.83
CV,%		13.3	7	2.3	17.9	9.4	11	12	18	29.35	14.32	26.34	18.62
LSD,0.05		0.7	9	1.0	11.0	7.9	5	7	10	0.23	0.22	0.44	0.75

<sup>1</sup> Vigor score based on a scale of 1 to 5 with 5 being the most vigorous seedling growth.

<sup>2</sup> Maturity rating scale: 37 = flag leaf emergence, 45 = boot swollen, 50 = beginning of inflorescence emergence, 58 = complete emergence of inflorescence, 62 = beginning of pollen shed. See Table 2 for complete scale.

\*Not significantly different from the highest numerical value in the column, based on the 0.05 LSD.

Nitrogen application: 60 lb/A of actual nitrogen on May 11 and August 7.

Rainfall deficit: May-August rainfall was 10.62 inches; rainfall deficit during this period in 2012 was -6.44 inches.

**Table 9. Dry matter yields, seedling vigor, percent stand, maturity and stand height of sorghum-sudangrass varieties sown May 28, 2013 at Lexington, Kentucky.**

Variety	Proprietor/ Distributor	Seedling Vigor <sup>1</sup> Jun 20	Percent Stand Jun 20	Maturity <sup>2</sup>			Height (inches)				Yield (DM tons/acre)				
				Jul 8	Aug 7	Sep 9	Jul 8	Aug 7	Sep 9	Oct 21	Jul 8	Aug 7	Sep 9	Oct 21	Total
<b>Commercial Varieties—Available for Farm Use</b>															
Greengrazer V	Farm Science Genetics	5.0	100	31.5	32.5	33.0	53	50	52	18	1.76	1.79	2.11	0.53	6.19*
Special Effort	Cisco	4.9	100	31.5	31.8	36.5	50	50	52	19	1.75	1.60	2.14	0.62	6.11*
SS211	Southern States	4.5	94	31.3	32.0	33.3	49	47	56	17	1.64	1.54	2.12	0.49	5.78*
NutraPlus BMR	Cisco	5.0	100	31.5	30.5	32.8	45	37	48	14	1.87	1.27	1.84	0.54	5.53*
SuperSugar	Gayland Ward Seed	4.8	97	31.8	32.5	33.5	50	52	51	19	1.59	1.53	1.82	0.50	5.45*
FSG214 BMR6	Farm Science Genetics	4.8	90	31.5	32.5	36.0	45	49	47	13	1.66	1.38	1.62	0.39	5.04
AS6503 BMR6	Alta Seeds/Ramer Seed	4.5	100	30.8	26.5	31.8	38	32	36	11	1.75	1.15	1.71	0.26	4.87
SweetSix BMR	Gayland Ward Seed	4.9	100	31.5	32.3	33.0	47	44	45	16	1.70	1.28	1.41	0.33	4.71
GW 300 BMR	Gayland Ward Seed	3.0	84	30.8	31.5	32.5	42	42	51	16	1.27	1.26	1.58	0.37	4.48
Sweet-for-Ever	Gayland Ward Seed	2.3	74	31.0	30.5	32.0	36	45	43	15	0.96	1.42	1.47	0.28	4.14
Sweet-for-Ever BMR	Gayland Ward Seed	3.5	90	30.5	30.0	31.3	36	36	35	12	1.23	0.99	1.13	0.21	3.56
Mean		4.3	93	31.2	31.1	31.2	44	44	47	15	1.56	1.38	1.72	0.41	5.08
CV,%		9.2	8	1.9	5.8	7.3	6	11	11	9	10.82	12.79	14.00	22.74	10.56
LSD,0.05		0.6	11	0.8	2.6	3.5	4	7	7	2	0.24	0.26	0.35	0.14	0.77

<sup>1</sup> Vigor score based on a scale of 1 to 5 with 5 being the most vigorous seedling growth.

<sup>2</sup> Maturity rating scale: 37 = flag leaf emergence, 45 = boot swollen, 50 = beginning of inflorescence emergence, 58 = complete emergence of inflorescence, 62 = beginning of pollen shed. See Table 2 for complete scale.

\*Not significantly different from the highest numerical value in the column, based on the 0.05 LSD.

Nitrogen application: 50 lb/A of actual nitrogen on July 22 and Aug 8.

**Table 10. Dry matter yields, stand rating, maturity and plant height of sorghum-sudangrass varieties sown May 21, 2014 at Lexington, Kentucky.**

Variety	Proprietor/ Distributor	Percent Stand Jun 17	Maturity <sup>1</sup>			Plant Height (in)			Yield (DM tons/acre)				
			Jul 8	Aug 13	Sep 16	Jul 8	Aug 13	Sep 16	Jul 8	Aug 13	Sep 16	Oct 29	Total
<b>Commercial Varieties—Available for Farm Use</b>													
FSG 214 BMR6	Farm Sci. Genetics/Allied	100	32.0	43.0	55.0	39	29	50	1.89	1.25	1.99	0.35	5.47*
GreenGrazer V	Farm Sci. Genetics/Allied	99	32.0	33.3	56.3	41	33	46	1.89	1.19	2.00	0.34	5.41*
NutraPlus BMR	Cisco	100	31.3	40.5	50.5	33	32	47	1.67	1.23	2.11	0.35	5.37*
Sordan Headless	Chromatin	100	31.0	31.8	37.0	35	32	45	1.49	1.28	2.29	0.26	5.32*
AS6503 BMR6	Atla Seeds/Ramer Seed	100	31.3	34.5	37.0	34	26	35	1.78	1.23	1.94	0.25	5.20*
SS211	Southern States	91	31.5	36.3	41.0	40	35	52	1.53	1.28	2.11	0.27	5.19*
Super Sugar (Delayed maturity)	Gayland Ward Seed	100	31.5	32.5	37.0	40	30	51	1.59	1.03	2.19	0.31	5.12*
SweetSix BMR	Gayland Ward Seed	100	31.3	34.8	49.0	35	29	48	1.50	1.29	1.97	0.34	5.10*
SDH2942 BMR	Chromatin	100	31.0	31.8	37.0	30	32	37	1.56	1.27	1.85	0.28	4.96*
Super Sugar Sterile	Gayland Ward Seed	99	32.3	33.3	51.0	38	32	47	1.55	1.05	1.87	0.28	4.76*
Special Effort	Cisco	100	31.3	39.5	50.8	33	32	44	1.45	1.06	1.73	0.38	4.62
GW 300 BMR	Gayland Ward Seed	98	31.5	31.0	41.0	38	26	44	1.56	0.82	1.64	0.07	4.09
<b>Experimental Varieties</b>													
ASBDSS	Farm Sci. Genetics/Allied	100	31.0	34.5	43.0	32	27	39	1.77	1.04	1.80	0.25	4.86*
SPX3952	Chromatin	100	31.0	41.8	45.0	29	29	34	1.29	1.14	1.53	0.36	4.32
Mean		99	31.40	35.6	45.0	35	30	44	1.63	1.15	1.93	0.29	4.99
CV,%		2	1.9	15.7	11.1	12	10	10	23.24	18.04	10.90	35.23	11.20
LSD,0.05		2	0.9	8.0	7.1	6	4	7	0.53	0.30	0.30	0.15	0.80

<sup>1</sup> Maturity rating scale: 37 = flag leaf emergence, 45 = boot swollen, 50 = beginning of inflorescence emergence, 58 = complete emergence of inflorescence, 62 = beginning of pollen shed. See Table 2 for complete scale.

\*Not significantly different from the highest numerical value in the column, based on the 0.05 LSD.  
Nitrogen application: 50 lb/A of actual nitrogen on May 27, July 11 and August 15.

**Table 11. Dry matter yields, seedling vigor, percent stand, maturity and stand height of teff varieties sown May 25, 2011 at Lexington, Kentucky.**

Variety <sup>3</sup>	Seedling Vigor <sup>1</sup> Jun 16	Percent Stand Jun 16	Maturity <sup>2</sup>		Height (inches) Jul 7	Yield (tons/acre)				
			Jul 7	Jul 22		Jul 7	Jul 22	Aug 15	Sep 27	Total
<b>Commercial Varieties—Available for Farm Use</b>										
Rooiberg	4.5	100	57.0	57.0	23	0.71	1.09	1.18	0.71	3.70*
Excaliber	4.0	100	55.0	56.0	25	0.62	1.02	1.12	0.89	3.65*
HorseCandi	4.0	99	47.5	51.3	21	0.71	0.99	1.06	0.88	3.64*
Pharaoh	4.9	100	44.5	53.5	23	0.78	0.97	1.03	0.78	3.56*
Witkope	4.0	100	55.5	56.0	24	0.69	1.11	0.97	0.70	3.47*
Corvallis	4.8	100	51.3	53.0	22	0.63	0.95	1.09	0.75	3.42*
Highveld	3.8	100	42.8	53.5	20	0.47	1.02	1.01	0.89	3.39*
Velvet	4.4	100	50.8	53.0	22	0.56	0.99	0.96	0.79	3.31*
Dessie	3.3	99	42.3	54.0	21	0.46	1.02	0.94	0.73	3.16*
Tiffany	4.0	100	46.5	54.5	19	0.41	1.00	0.96	0.78	3.14*
VA-T1Brown	4.8	100	48.0	52.0	20	0.45	0.95	1.00	0.68	3.07
Summer Delight	3.3	99	48.8	54.0	17	0.44	0.93	0.91	0.70	2.98
Mean	4.1	100	49.1	54.0	21	0.58	1.00	1.02	0.77	3.37
CV,%	18.4	1	9.9	3.4	13	46.81	8.95	11.59	16.40	12.33
LSD,0.05	1.1	1	7.0	2.6	4	0.39	0.13	0.17	0.18	0.60

<sup>1</sup> Vigor score based on a scale of 1 to 5 with 5 being the most vigorous seedling growth.

<sup>2</sup> Maturity rating scale: 37 = flag leaf emergence, 45 = boot swollen, 50 = beginning of inflorescence emergence, 58 = complete emergence of inflorescence, 62 = beginning of pollen shed. See Table 2 for complete scale.

<sup>3</sup> Check with local dealer for available varieties.

\*Not significantly different from the highest numerical value in the column, based on the 0.05 LSD.  
Nitrogen application: 30 lb/A on June 2, 40 lb/A on July 7 and July 27 and 30 lb/A on Aug 19 of actual nitrogen.

**Table 12. Dry matter yields, seedling vigor, percent stand and maturity of teff varieties sown May 10, 2012 at Lexington, Kentucky.**

Variety <sup>3</sup>	Seedling Vigor <sup>1</sup> Jun 4	Percent Stand Jun 4	Maturity <sup>2</sup>			Yield (tons/acre)				
			Jun 27	Jul 25	Aug 10	Jun 27	Jul 25	Aug 10	Sept 27	Total
<b>Commercial Varieties—Available for Farm Use</b>										
Rooiberg	4.0	100	57.0	60.0	57.5	0.90	0.40	0.84	1.08	3.21*
Highveld	4.1	99	52.5	59.0	55.0	0.77	0.41	0.77	1.11	3.05*
Excaliber	4.1	100	54.5	59.0	56.5	0.77	0.38	0.72	1.08	2.95*
Tiffany	4.3	100	48.5	56.5	52.0	0.74	0.34	0.73	1.05	2.85*
Witkope	3.8	99	56.0	57.0	56.0	0.68	0.34	0.75	1.04	2.82*
Pharaoh	4.1	100	46.3	52.8	49.8	0.64	0.29	0.73	1.05	2.71
Corvalis	4.4	100	47.5	56.0	51.3	0.63	0.34	0.72	1.00	2.68
Dessie	3.6	100	52.0	57.0	54.0	0.56	0.34	0.70	1.05	2.66
Velvet	3.9	100	53.5	58.0	54.0	0.58	0.32	0.74	1.01	2.65
VA-T1Brown	4.4	100	50.3	53.0	46.8	0.58	0.35	0.71	1.00	2.63
HorseCandi	3.9	100	48.0	55.0	50.3	0.60	0.27	0.71	1.04	2.63
SummerDelight	4.1	100	51.5	57.5	51.0	0.54	0.29	0.72	1.04	2.60
<b>Experimental Varieties</b>										
F-11	3.5	99	46.8	55.5	54.0	0.53	0.26	0.64	0.90	2.33
Mean	4.0	100	51.1	56.6	52.9	0.66	0.33	0.73	1.04	2.75
CV,%	19.8	1	5.4	3.6	4.1	29.04	21.35	11.73	10.51	12.07
LSD,0.05	1.1	1	3.9	2.9	3.1	0.27	0.10	0.12	0.16	0.48

<sup>1</sup> Vigor score based on a scale of 1 to 5 with 5 being the most vigorous seedling growth.

<sup>2</sup> Maturity rating scale: 37 = flag leaf emergence, 45 = boot swollen, 50 = beginning of inflorescence emergence, 58 = complete emergence of inflorescence, 62 = beginning of pollen shed. See Table 2 for complete scale.

<sup>3</sup> Check with local dealer for available varieties.

\*Not significantly different from the highest numerical value in the column, based on the 0.05 LSD.

Nitrogen application: 40 lb/A on May 11, 50 lb/A on July 26 and 30 lb/A on August 14 of actual nitrogen.

Rainfall deficit: May-August rainfall was 10.62 inches; rainfall deficit during this period in 2012 was -6.44 inches.

**Table 13. Dry matter yields, seedling vigor, percent stand and maturity of teff varieties sown May 28, 2013 at Lexington, Kentucky.**

Variety <sup>3</sup>	Seedling Vigor <sup>1</sup> Jun 20	Percent Stand Jun 20	Maturity <sup>2</sup>		Yield (tons/acre)				Total
			Jul 17	Aug 7	Jul 17	Aug 7	Sep 19	Oct 21	
<b>Commercial Varieties—Available for Farm Use</b>									
Witkope	2.4	93	56.0	49.0	1.98	1.20	1.12	0.61	4.90*
Excaliber	3.4	99	52.5	51.5	1.99	1.17	1.18	0.54	4.87*
Highveld	3.1	98	49.8	32.0	1.94	0.96	1.47	0.48	4.86*
Velvet	2.6	98	53.0	40.8	2.15	1.04	1.11	0.55	4.84*
Rooiberg	2.8	97	56.0	48.8	2.09	1.07	1.24	0.44	4.83*
Pharaoh	3.3	99	45.0	35.3	2.03	1.14	1.09	0.52	4.78*
Corvalis	2.5	98	48.0	38.5	1.95	1.09	1.18	0.51	4.73*
SummerDelight	3.9	99	48.0	35.3	2.14	1.04	1.04	0.51	4.72*
VA-T1Brown	2.9	99	51.3	37.0	2.10	1.03	1.08	0.38	4.60*
Tiffany	2.9	100	49.0	32.0	1.95	1.08	1.02	0.55	4.60*
Dessie	2.6	95	54.0	43.0	1.88	1.04	1.17	0.49	4.59*
HorseCandi	2.0	95	49.8	40.3	2.01	1.09	0.98	0.47	4.56*
Moxie	2.6	97	53.5	33.8	2.04	0.93	1.02	0.45	4.45*
<b>Experimental Varieties</b>									
F11	2.8	100	46.3	37.0	2.08	1.04	1.00	0.56	4.67*
Mean	2.8	98	50.9	39.6	2.02	1.07	1.12	0.50	4.21
CV,%	43.0	4	6.0	16.3	8.12	13.47	19.82	20.51	9.02
LSD,0.05	1.7	6	4.3	9.2	0.23	0.21	0.32	0.15	0.61

<sup>1</sup> Vigor score based on a scale of 1 to 5 with 5 being the most vigorous seedling growth.

<sup>2</sup> Maturity rating scale: 37 = flag leaf emergence, 45 = boot swollen, 50 = beginning of inflorescence emergence, 58 = complete emergence of inflorescence, 62 = beginning of pollen shed. See Table 2 for complete scale.

<sup>3</sup> Check with local dealers for available varieties.

\*Not significantly different from the highest numerical value in the column, based on the 0.05 LSD.

Nitrogen application: 40 lb/A on June 5 and 50 lb/A on July 22 and Aug 8 of actual nitrogen.

**Table 14. Dry matter yields, stand rating and maturity of teff varieties sown May 21, 2014 at Lexington, Kentucky.**

Variety <sup>2</sup>	Percent Stand Jun 17	Maturity <sup>1</sup>		Yield (tons/acre)				
		Jul 8	Aug 4	Jul 8	Aug 4	Aug 29	Oct 13	Total
<b>Commercial Varieties—Available for Farm Use</b>								
Summer Delight	100	55.5	54.0	1.63	1.01	1.78	0.93	5.34*
Corvalis	100	52.3	52.5	1.27	1.05	1.64	0.98	4.95*
Witkope	96	56.0	59.0	1.09	1.09	1.68	0.94	4.81*
VA-T1Brown	100	55.5	52.0	1.15	1.09	1.70	0.76	4.70*
Tiffany	98	54.0	53.0	1.23	1.17	1.55	0.73	4.69*
Highveld	89	50.5	54.5	0.89	1.11	1.74	0.85	4.59*
Dessie	95	54.0	56.0	0.97	1.18	1.39	1.00	4.55*
Moxie	100	52.8	54.0	1.28	1.11	1.40	0.53	4.33*
Velvet	100	56.0	54.0	0.78	0.91	1.60	1.00	4.29*
Pharaoh	97	51.7	51.5	1.09	0.91	1.42	0.75	4.18*
Rooiberg	98	54.5	59.0	0.66	1.05	1.44	0.82	3.97*
HorseCandi	98	54.5	53.0	0.78	0.89	1.30	0.61	3.58
<b>Experimental Varieties</b>								
F11	99	50.5	53.5	0.97	0.89	1.30	0.87	4.04*
Mean	98	53.7	54.3	1.06	1.04	1.54	0.83	4.47
CV,%	6	6.8	4.3	44.66	16.00	23.91	38.16	20.95
LSD,0.05	8	5.5	3.4	0.70	0.24	0.54	0.46	1.38

<sup>1</sup> Maturity rating scale: 37 = flag leaf emergence, 45 = boot swollen, 50 = beginning of inflorescence emergence, 58 = complete emergence of inflorescence, 62 = beginning of pollen shed. See Table 2 for complete scale.

<sup>2</sup> Check with local dealer for available varieties.

\*Not significantly different from the highest numerical value in the column, based on the 0.05 LSD. Nitrogen application: 50 lb/A of actual nitrogen on May 27 and July 11.

**Table 15. Dry matter yields, seedling vigor, percent stand and maturity of pearl millet varieties sown May 28, 2013 at Lexington, Kentucky.**

Variety	Proprietor/Distributor	Seedling Vigor <sup>1</sup> Jun 20	Percent Stand Jun 20	Maturity <sup>2</sup>			Plant Height (inches)				Yield (tons/acre)				
				Jul 12	Aug 7	Sep 9	Jul 12	Aug 7	Sep 9	Oct 21	Jul 15	Aug 7	Sep 9	Oct 21	Total
<b>Commercial Varieties—Available for Farm Use</b>															
Tifleaf III Hybrid	Gayland Ward Seed	4.8	99	30.0	48.5	64	32	35	45	21	1.64	1.65	2.26	0.62	6.16*
SS635	Southern States	3.5	89	29.5	37.0	64	32	32	47	22	1.61	1.39	2.19	0.55	5.74*
Pennleaf Hybrid	Pennington Seed	2.5	74	30.0	46.3	64	30	34	43	19	1.47	1.29	1.75	0.44	4.95*
PP102M Hybrid	Cisco	2.3	74	29.5	46.3	64	34	33	53	19	1.59	1.16	1.84	0.35	4.93*
SS501	Southern States	3.0	76	30.0	33.3	64	42	32	59	21	1.57	1.08	1.84	0.31	4.80*
Mean		3.2	82	29.8	42.3	64	34	33	49	21	1.58	1.31	1.97	0.45	5.32
CV,%		32.7	10	4.1	19.7	0	11	6	9	15	25.54	11.75	16.09	23.58	17.04
LSD,0.05		1.6	13	1.9	12.8	0	6	3	7	5	0.62	0.24	0.49	0.23	1.40

<sup>1</sup> Vigor score based on a scale of 1 to 5 with 5 being the most vigorous seedling growth.

<sup>2</sup> Maturity rating scale: 37 = flag leaf emergence, 45 = boot swollen, 50 = beginning of inflorescence emergence, 58 = complete emergence of inflorescence, 62 = beginning of pollen shed. See Table 2 for complete scale.

\*Not significantly different from the highest numerical value in the column, based on the 0.05 LSD. Nitrogen application: 50 lb/A of actual nitrogen on July 22 and Aug 8.



**Table 16. Dry matter yields, stand rating, maturity and plant height of pearl millet varieties sown May 21, 2014 at Lexington, Kentucky.**

Variety	Proprietor/ Distributor	Percent Stand Jun 17	Maturity <sup>1</sup>		Plant Height (in)			Yield (tons/acre)				
			Jul 15	Aug 11	Jul 15	Aug 11	Sep 12	Jul 15	Aug 11	Sep 12	Oct 29	Total
<b>Commercial Varieties—Available for Farm Use</b>												
SS635	Southern States	97	31.0	60.0	30	34	38	1.41	1.55	1.72	0.56	5.24*
Tifleaf III Hybrid	Gayland Ward Seed	99	31.5	61.0	27	35	35	1.26	1.27	1.74	0.67	4.94*
SS501	Southern States	97	31.8	46.3	38	29	47	1.75	0.91	1.69	0.28	4.64*
PP102M Hybrid	Cisco	100	32.0	55.0	34	27	42	1.63	0.95	1.48	0.27	4.33*
Pennleaf Hybrid	Pennington Seed	98	31.0	58.0	27	28	33	1.21	1.04	1.57	0.43	4.25*
Mean		98	31.5	56.1	31	30	39	1.45	1.15	1.64	0.44	4.68
CV,%		3	2.2	4.1	7	9	12	12.19	21.54	17.91	24.97	13.69
LSD,0.05		4	1.1	3.5	3	4	7	0.27	0.38	0.45	0.17	0.99

<sup>1</sup> Maturity rating scale: 37 = flag leaf emergence, 45 = boot swollen, 50 = beginning of inflorescence emergence, 58 = complete emergence of inflorescence, 62 = beginning of pollen shed. See Table 2 for complete scale.

\*Not significantly different from the highest numerical value in the column, based on the 0.05 LSD.

Nitrogen application: 50 lb/A of actual nitrogen on May 27, July 21 and August 15.

**Table 17. Dry matter yields, seedling vigor, percent stand, heading date, harvest height and harvest date of forage sorghum varieties sown May 28, 2013 at Lexington, Kentucky.**

Variety	Proprietor/Distributor	Seedling Vigor <sup>1</sup> Jun 20	Percent Stand Jun 20	Heading Date <sup>2</sup>	Harvest Height (feet)	Harvest Date <sup>3</sup>	Yield (DM tons/ acre)
<b>Commercial Varieties—Available for Farm Use</b>							
Ensilemaster	Caudill Seed	2.6	81	Aug 22	10.6	Sept 10	10.41*
GW2120	Gayland Ward Seed	3.5	91	Aug 13	9.4	Sept 6	9.77*
GW400 BMR	Gayland Ward Seed	3.5	87	Aug 13	9.9	Sept 6	7.81
AF7201 BMR6	Alta Seeds/Ramer Seed	4.0	90	Aug 9	9.8	Aug 27	7.45
AF7401 BMR6	Alta Seeds/Ramer Seed	4.9	98	Aug 21	6.5	Sept 10	6.36
<b>Experimental Varieties</b>							
Exp10074	Gayland Ward Seed	4.9	97	Aug 2	10.6	Aug 27	8.49
Mean		3.9	91		9.5		8.38
CV,%		11.4	6		3.0		12.63
LSD,0.05		0.7	8		0.4		1.60

<sup>1</sup> Vigor score based on a scale of 1 to 5 with 5 being the most vigorous seedling growth.

<sup>2</sup> Approximately 50% of heads fully emerged.

<sup>3</sup> Harvested at soft dough stage.

\*Not significantly different from the highest numerical value in the column, based on the 0.05 LSD.

**Table 18. Dry matter yields, stand rating, heading date, plant height and maturity of forage sorghum varieties sown May 21, 2014 at Lexington, Kentucky.**

Variety	Proprietor/Distributor	Percent Stand Jun 17	Heading Date <sup>1</sup>	Plant Height (ft) Sep 18	Maturity <sup>2</sup>	Yield (DM tons/acre)	Forage Quality Analysis <sup>3</sup>				
							%CP	%ADF	%NDF	%TDN	RFV
SS405	Chromatin	98	Aug 25	9.9	86.5	11.88*	3.4	36.5	58	57	97
SD1741 BMR	Chromatin	97	Jul 29	8.5	93.0	8.39	7.8	35.6	59.2	57	96
NK300	Chromatin	97	Aug 12	6.0	92.5	7.94	6.1	32.9	51.6	57	114
1990	Chromatin	88	—	9.9	—	7.62	4.7	43.2	70.3	54	73
GW600 BMR	Gayland Ward Seed	94	Aug 11	8.3	92.0	6.79	8.2	27.8	46.4	60	135
FSG114 BMR6	Farm Science Genetics	93	Aug 5	6.6	90.5	5.96	5.5	34	54	58	108
AF7401 BMR6	Alta Seeds/Ramer Seed	100	Aug 19	5.3	89.5	5.96	6.3	28.5	47.7	59	130
EnsilMaster	Caudill Seed	59 <sup>4</sup>	Aug 26	8.5	84.5	5.66	3.4	33.9	59.6	57	104
GW2120	Gayland Ward Seed	93	Aug 6	6.9	91.0	5.60	6.2	29.6	48.7	59	126
AF7201 BMR6	Alta Seeds/Ramer Seed	90	Aug 10	7.0	91.0	5.12	2.5	39.1	66.5	54	82
GW400 BMR	Gayland Ward Seed	94	Aug 4	7.1	89.5	4.99	6.2	28.5	48.8	59	127
FSG115Dwarf BMR6	Farm Science Genetics	53 <sup>4</sup>	Aug 26	6.4	92.5	3.24	7.5	36.2	57.8	57	97
SilageKingDwarf BMR6	Gayland Ward Seed	39 <sup>4</sup>	Aug 25	6.1	91.0	3.02	8	32.1	55	58	108
<b>Experimental Varieties</b>											
SPX28313	Chromatin	93	Aug 27	11.0	77.5	13.28*	4.5	34.5	58.8	57	98
SPX902	Chromatin	98	—	10.9	—	11.47	3.5	41.8	64.7	55	81
SPX904	Chromatin	99	—	10.9	—	11.07	5.6	40.1	68.5	54	78
SPX903	Chromatin	98	—	11.9	—	9.78	4.2	42.3	69.1	54	75
SPX901	Chromatin	92	—	10.8	—	7.89	7.7	40.5	68	54	78
X942BMR	Chromatin	99	—	9.0	—	7.38	7.6	34.1	59.3	57	98
SPX3952	Chromatin	98	Aug 1	7.8	92.5	5.19	6.7	35.4	59	57	97
SPX3902	Chromatin	88	Aug 22	6.1	89.5	4.34	6.3	32.2	55.2	58	108
SPX3903	Chromatin	93	Aug 24	5.1	92.5	3.43	9.5	31.1	55.7	58	108
Mean		89	Aug 14	8.2	89.7	7.09					
CV,%		10	4 days	12.6	3.5	17.35					
LSD,0.05		13	5 days	1.5	4.5	1.74					

<sup>1</sup> Approximately 50% of heads fully emerged. Those without a date are photoperiod sensitive.

<sup>2</sup> See Table 2 for maturity scale.

<sup>3</sup> CP = crude protein, ADF = acid detergent fiber, NDF = neutral detergent fiber, TDN = total digestible nutrients, RFV = relative feed value. All were harvested on the same day, therefore quality may not be optimal for some varieties.

<sup>4</sup> Poor germination of these varieties resulted in reduced stand and yield.

\*Not significantly different from the highest numerical value in the column, based on the 0.05 LSD.

Nitrogen application: 60 lb/A of actual nitrogen on May 27.

**Table 19. Summary of Kentucky sudangrass yield trials 2008-2014 (yield shown as a percentage of the mean of the commercial varieties in the trial).**

Variety	Proprietor/KY Distributor	Lexington							Mean <sup>3</sup> (#trials)
		2008 <sup>1,2</sup>	2009	2010	2011	2012	2013	2014	
AS9301 BMR <sup>4</sup>	Alta Seeds/Ramer Seed					118			—
Enorma BMR	Cal/West Seeds			99	94	92	91	83	92(5)
Hayking BMR	Central Farm Supply	111	112	91	97	97	96	92	99(7)
Monarch V	Public	104	96	102	97	93	98	110	100(7)
Piper	Public	90	91	97	94	104	105	89	96(7)
ProMax BMR	Ampac Seed	95	101	110	115	96	103	100	103(7)
SS130 BMR	Cal/West Seeds			101	103		107	106	104(4)
Trudan Headless	Chromatin							118	—

<sup>1</sup> Establishment year.

<sup>2</sup> Use this summary table as a guide in making variety decisions, but refer to specific tables in this report to determine statistical differences in forage yield between varieties.

<sup>3</sup> Mean only presented when respective variety was included in two or more trials.

<sup>4</sup> BMR (Brown Mid-rib) means that a variety has been developed to produce lower amounts of lignin which usually translates into higher quality.

**Table 20. Summary of Kentucky sorghum-sudangrass yield trials 2008-2014 (yield shown as a percentage of the mean of the commercial varieties in the trial).**

Variety	Proprietor/KY Distributor	Lexington							Mean <sup>3</sup> (#trials)
		2008 <sup>1,2</sup>	2009	2010	2011	2012	2013	2014	
		All trials are 1 year yields							
AS6402 BMR <sup>4</sup>	Alta Seeds/Ramer Seed					91			–
AS6503 BMR6	Alta Seeds/Ramer Seed						96	103	100(2)
FSG 208 BMR	Farm Science Genetics			75					–
FSG 214 BMR6	Farm Science Genetics					99	108		104(2)
Greengrazer V	Farm Science Genetics			166			122	107	131(3)
GW300 BMR	Gayland Ward Seed				88	78	88	81	84(4)
HyGain	Turner Seed	104	105	118					109(3)
MS 202 BMR	Farm Science Genetics			106					–
NutraPlus BMR	Cisco	106	97	94	103	106	109	106	103(7)
Sordan Headless	Chromatin							105	–
Special Effort	Cisco	109	110	93	94	115	120	91	105(7)
SS211	Southern States				104	93	114	103	104(4)
SS220 BMR	Southern States		107	84		112			101(3)
Surpass BMR-6	Turner Seed	81	80	64					75(3)
Super Sugar	Gayland Ward Seed				102	117	107		109(3)
Super Sugar Delayed maturity	Gayland Ward Seed							101	–
Super Sugar Sterile	Gayland Ward Seed							94	–
Sweet-For-Ever	Gayland Ward Seed				110	107	81		99(3)
Sweet-For-Ever BMR	Gayland Ward Seed					78	70		74(2)
SweetSix BMR	Gayland Ward Seed						93	101	97(2)
Vita-Cane	Gayland Ward Seed					121			–

<sup>1</sup> Establishment year.

<sup>2</sup> Use this summary table as a guide in making variety decisions, but refer to specific tables in this report to determine statistical differences in forage yield between varieties.

<sup>3</sup> Mean only presented when respective variety was included in two or more trials.

<sup>4</sup> BMR (Brown Mid-rib) means that a variety has been developed to produce lower amounts of lignin which usually translates into higher quality.

**Table 21. Summary of Kentucky teff yield trials 2008-2014 (yield shown as a percentage of the mean of the commercial varieties in the trial).**

Variety	Princeton		Lexington							Mean <sup>3</sup> (#trials)
	2008 <sup>1,2</sup>	2009	2008	2009	2010	2011	2012	2013	2014	
	All trials are 1 year yields									
Corvallis	94	112	81	101	91	101	96	100	110	98(9)
Dessie	102	87	99	92	96	94	95	97	101	96(9)
Excaliber	109	111	109	104	125	108	106	103		109(8)
Highveld	111	115	100	121	106	101	109	103	102	108(9)
HorseCandi	91	84	99	105	89	108	94	97	80	94(9)
Moxie								94	96	95(2)
Pharaoh	95	101	105	85	106	106	97	101	93	99(9)
Rooiberg	102	107	112	109	113	108	115	102	88	106(9)
Summer Delight		90		91	96	88	93	100	119	97(7)
Tiffany	102	106	102	93	82	93	102	98	104	98(9)
VA T1 Brown		89		99	87	91	94	98	104	95(7)
Velvet		94		100	97	98	95	103	95	97(7)
Witkope	94	100	93	101	115	103	101	104	107	102(9)

<sup>1</sup> Establishment year.

<sup>2</sup> Use this summary table as a guide in making variety decisions, but refer to specific tables in this report to determine statistical differences in forage yield between varieties.

<sup>3</sup> Mean only presented when respective variety was included in two or more trials.

**Table 22. Summary of Kentucky pearl millet yield trials 2012-2014 (yield shown as a percentage of the mean of the commercial varieties in the trial).**

Variety	Proprietor/KY Distributor	Lexington			Mean <sup>3</sup> (#trials)
		2012 <sup>1,2</sup>	2013	2014	
Pennleaf Hybrid	Pennington Seed	117	93	91	100(3)
PP102M Hybrid	Cisco	93	93	93	93(3)
SS501	Southern States		90	99	95(2)
SS635	Southern States		108	112	110(2)
Tiffleaf III Hybrid	Gayland Ward Seed	90	116	106	104(3)

<sup>1</sup> Establishment year.

<sup>2</sup> Use this summary table as a guide in making variety decisions, but refer to specific tables in this report to determine statistical differences in forage yield between varieties.

<sup>3</sup> Mean only presented when respective variety was included in two or more trials.



Mention or display of a trademark, proprietary product, or firm in text or figures does not constitute an endorsement and does not imply approval to the exclusion of other suitable products or firms.