

# 2006 Alfalfa Grazing Tolerance Report

G.L. Olson, S.R. Smith, G.D. Lacefield, and E. Vanzant

## Introduction

Alfalfa (*Medicago sativa*) is the highest yielding, highest quality forage legume grown in Kentucky. It forms the basis of Kentucky's cash hay enterprise and is an important component in dairy, horse, beef, and sheep diets. Recent emphasis on its use as a grazing crop and the release of grazing tolerant varieties have raised the following question: Do varieties differ in tolerance to grazing? We have chosen to use the standard tolerance test recommended by the North American Alfalfa Improvement Conference. This test uses continuous heavy grazing to sort out differences in grazing tolerance in a relatively short period of time.

This report summarizes current research on the grazing tolerance of alfalfa varieties when subjected to continuous heavy grazing pressure during the grazing season. New for 2006, Table 5 shows a summary of all alfalfa varieties tested in Kentucky during the last 12 years. Go to the UK Forage Extension Web site at <[www.uky.edu/Ag/Forage](http://www.uky.edu/Ag/Forage)> to obtain electronic versions of all forage variety testing reports from Kentucky, from surrounding states and a large number of other forage publications.

## Description of the Tests

Alfalfa variety tests for grazing tolerance were established in Lexington in the fall of 2004 and 2005. The soils at this location are well-drained silt loams and are well suited to alfalfa. Plots were 5 by 15 feet in a randomized complete block design, with each variety replicated six times. In each test, 20 pounds of seed per acre were planted into a prepared seedbed using a

disk drill. All seed lots were treated with metalaxyl fungicide and inoculated if not supplied with these treatments. Plots were grazed continuously beginning the first spring after seeding. Grazing pressure was maintained to keep plant height to less than 3 inches. In general, plots were grazed from April until mid-September. Supplemental hay was fed during periods of slowest growth. To check stand survival after the grazing season, visual ratings of percent stand were made in the fall several weeks after the cattle were removed and in the spring prior to grazing to check on winter survival and spring growth. Since trials were seeded in rows, persistence ratings were based on density within a row and not total ground cover. Pests (weeds and insects) were controlled so they would not limit yield or persistence. Fertilizers (lime, P, K, and Boron) were applied as needed. In each trial, Alfagrade was the grazing-tolerant check variety, and either Apollo or 5432 was the grazing-intolerant check variety.

## Results and Discussion

Weather data for Lexington for 2004, 2005 and 2006 are presented in Table 1.

Data on percent stand are presented in Tables 2 and 3. Statistical analyses were performed on all alfalfa yield data (including experimentals) to determine if the apparent differences are truly due to variety or just due to chance. Varieties not significantly different from the highest numerical value in a column are marked with one asterisk (\*). To determine if two varieties are truly different, compare the difference between the two varieties

**Table 1. Temperature and rainfall at Lexington, Kentucky, in 2002, 2003, 2004, 2005, and 2006.**

	2002				2003				2004				2005				2006			
	Temp.		Rainfall		Temp.		Rainfall		Temp.		Rainfall		Temp.		Rainfall		Temp.		Rainfall	
	°F	DEP	IN	DEP	°F	DEP	IN	DEP	°F	DEP	IN	DEP	°F	DEP	IN	DEP	°F	DEP	IN	DEP
JAN	38	+7	2.12	-0.74	26	-5	0.96	-1.90	30	-1	3.14	+0.28	37	+6	4.35	+1.49	42	+11	4.77	+1.91
FEB	38	+3	1.28	-1.93	32	-3	3.59	+0.38	36	+1	1.32	-1.89	39	+4	1.68	-1.53	36	+1	2.13	-1.08
MAR	45	+1	7.93	+3.53	47	+3	2.09	-2.31	47	+3	3.43	-0.97	41	-3	2.79	-1.61	44	0	3.05	-1.35
APR	58	+3	4.19	+0.31	57	+2	3.14	-0.74	55	0	3.06	-0.82	56	+1	3.30	-0.58	59	+4	3.52	-0.36
MAY	61	-3	4.36	-0.11	63	-1	6.68	+2.21	68	+4	9.79	+5.32	61	-3	1.78	-2.69	62	-2	2.99	-1.48
JUN	74	+2	2.45	-1.21	69	-3	4.85	+1.19	72	0	3.13	-0.53	75	+3	1.33	-2.33	70	-2	1.82	-1.84
JUL	78	+2	1.10	-3.90	74	-2	2.68	-2.32	73	-3	7.65	+2.65	77	+1	3.30	-1.70	76	0	5.13	+0.13
AUG	77	+2	0.95	-2.98	75	0	5.26	+1.33	71	-4	2.91	-1.02	78	+3	3.34	-0.59	76	+1	3.23	-0.70
SEP	72	+4	4.90	+1.70	65	-3	4.22	+1.02	68	0	2.61	-0.59	72	+4	0.59	-2.21	64	-4	9.27	+6.07
OCT	55	-2	5.61	+3.04	56	-1	1.61	-0.96	58	+1	5.65	+3.08	58	+1	0.92	-1.65	54	-3	4.88	+2.31
NOV	43	-2	3.76	+0.37	50	+5	4.63	+1.24	49	+4	6.29	+2.90	47	+2	1.54	-1.85	47	+2	1.78	-1.61
DEC	36	0	4.11	-1.13	36	0	3.26	-0.72	36	0	3.20	-0.78	32	-4	2.19	-1.79				
Total			42.73	-1.79			42.97	-1.58			52.18	+7.63			27.51	-17.04			42.57	+2.00

DEP is departure from the long-term average.

to the Least Significant Difference (LSD) at the bottom of the column. If the difference is equal to or greater than the LSD, the varieties are truly different when grown under the conditions at a given location. The Coefficient of Variation (CV), which is a measure of the variability of the data, is included for each column of means. Low variability is desirable, and increased variability within a study results in higher CVs and larger LSDs.

Apollo and 5432 have been used widely in trials as the grazing-intolerant varieties. Therefore, the response of these varieties provides a useful measure of the severity of the grazing pressure applied to the plots. In general, types developed for tolerance to grazing tolerated heavy grazing pressure better than hay types. Table 4 summarizes information about distributors, fall dormancy ratings, disease resistance information, and persistence across years for all varieties included in these tests.

Table 5 is a summary of stand persistence data from 1994-2006 of commercial varieties that have been entered in the Kentucky trials. The data for each specific trial is listed as a percentage of the grazing tolerant variety Aflagraze. In other words, in each trial Aflagraze is 100%—varieties with percentages over 100 persisted better than Aflagraze and varieties with percentages less than 100 persisted less than Aflagraze. Direct, statistical comparisons of varieties cannot be made using the summary Table 5, but these comparisons do help to identify varieties for further consideration. Varieties that have performed better than average over many years and at several locations have very stable performance, while others may have performed very well in wet years or on particular soil types. These details may influence variety choice and the information can be found in the yearly reports. See footnote in Table 5 to determine which yearly report to refer to.

**Table 2. Percent stand and seedling vigor rating of alfalfa varieties sown Sept. 3, 2004, in a cattle grazing tolerance study at Lexington, Kentucky.**

Variety	Seedling Vigor <sup>1</sup> Nov 8, 2004	Percent Stand			
		Apr 8, 2005	Oct 31, 2005	Apr 6, 2006	Oct 23, 2006
<b>Commercial Varieties—Available for Farm Use</b>					
Aflagraze	3.3	83	97	98	45*
5432	3.2	88	95	97	25
<b>Experimental Varieties</b>					
GA984	4.3	80	98	100	50*
GA1-01-1	3.5	76	95	95	47*
GA4-01-1	4.7	86	98	100	45*
GA3-01-1	4.2	83	97	98	42*
Mean	3.9	83	97	98	42
CV,%	17.5	13	4	6	32
LSD,0.05	0.8	13	5	7	16
*Not significantly different from the highest numerical value in the column, based on the 0.05 LSD.					
<sup>1</sup> Vigor score based on a scale of 1 to 5 with 5 being the most vigorous seedling growth.					

## Summary

Measurements taken after multiple years of grazing in these trials indicate that alfalfa varieties have been developed that exhibit improved tolerance to heavy continuous grazing pressure compared to standard hay-type varieties. The grazing management imposed in these trials included continuous stocking from the initiation of grazing in spring until mid-September, when grazing was terminated for the season to allow stands to acclimate to winter. Heavy grazing pressure was used purposely in these trials to better differentiate among varieties for relative grazing tolerance. Research has shown that abusive grazing tests are a good way to sort out differences in grazing tolerance between varieties in a relatively short period of time. Recommended rotational grazing management would improve alfalfa forage productivity and stand persistence.

The information in this report should be used in conjunction with other yield, pest resistance, and adaptation information in selecting the best alfalfa varieties for use in each individual situation.

Good management for maximum life when grazing alfalfa includes:

- allowing grazing alfalfa to become completely established before grazing.
- using rotational grazing where animals harvest available forage in seven days or less, followed by resting for 28 days before regazing.
- adding any needed fertilizer and lime.
- removing grazing livestock from alfalfa fields from mid-September until Nov. 1 to replenish root reserves for winter survival.

**Table 3. Percent stand and seedling vigor rating of alfalfa varieties sown Sept. 8, 2005, in a cattle grazing tolerance study at Lexington, Kentucky.**

Variety	Seedling Vigor <sup>1</sup> Nov 7, 2005	Percent Stand	
		Apr 17, 2006	Oct 20, 2006
Ameristand 407TQ	3.2	88	88*
Integrity	3.2	85	88*
Ameristand 403T	3.2	83	87*
Triple Trust 450	3.7	79	83*
Spredor 3	3.5	83	78*
Apollo	3.3	77	72
Aflagraze	2.3	78	68
Mean	3.2	82	80
CV,%	33.6	11	11
LSD,0.05	1.3	10	11
*Not significantly different from the highest numerical value in the column, based on the 0.05 LSD.			
<sup>1</sup> Vigor score based on a scale of 1 to 5 with 5 being the most vigorous seedling growth.			

**Table 4. Characterization and summary of persistence of alfalfa varieties under heavy grazing pressure across years at Lexington.**

Variety/Proprietor		Variety Characteristics <sup>1</sup>						2004 <sup>3</sup>				2005	
		FD <sup>4</sup>	Disease Resistance <sup>2</sup>					Apr 2005 <sup>5</sup>	Oct 2005	Apr 2006	Oct 2006	Apr 2006	Oct 2006
			BW	FW	AN	PRR	APH						
<b>Commercial Varieties—Available for Farm Use</b>													
Alfagraze	America's Alfalfa	4	MR	R	MR	LR	-	*	*	*	*	*	X
Ameristand 403T	America's Alfalfa	4	HR	HR	HR	HR	R					*	*
Ameristand 407TQ	America's Alfalfa											*	*
Apollo	ABI/America's Alfalfa	4	R	R	LR	R	-					X	X
Integrity	PGI Alfalfa, Inc.	4	HR	HR	HR	HR	HR					*	*
5432	Pioneer	4	HR	HR	-	MR	-	*	*	*	X		
Spredor 3	Syngenta	1	HR	HR	R	MR	S					*	*
Triple Trust 450	ABI/America's Alfalfa	5	HR	HR	HR	HR	HR					*	*
<b>Experimental Varieties</b>													
GA1-01-1	University of Georgia							*	*	*	*		
GA3-01-1	University of Georgia							*	*	*	*		
GA4-01-1	University of Georgia							*	*	*	*		
GA984	University of Georgia							*	*	*	*		
<sup>1</sup> Variety Characteristics: FD=Fall Dormancy, BW=Bacterial Wilt, FW=Fusarium Wilt, AN=Anthracnose, PRR=Phytophthora Root Rot, APH=Aphanomyces Root Rot. <sup>2</sup> Disease Resistance: S=Susceptible, LR=Low Resistance, MR=Medium Resistance, R=Resistance, HR=High Resistance. <sup>3</sup> Establishment year. <sup>4</sup> Fall Dormancy: 2=Vernal, 3=Ranger, 4=Saranac, 5=DuPuits. <sup>5</sup> Date of rating percent stand. *Not significantly different from the most persistent variety. An "x" in the block indicates the variety was in the test but the stand survival was significantly less than the most persistent variety. An open block indicates the variety was not in the test.													

**Authors:**

- G.L. Olson, Research Specialist, Forages, UK Department of Plant and Soil Sciences
- S.R. Smith, Extension Associate Professor, Forages, UK Department of Plant and Soil Sciences
- G.D. Lacefield, Extension Professor, Forages, UK Department of Plant and Soil Sciences
- E. Vanzant, Associate Professor, Beef Cattle Nutrition, UK Department of Animal and Food Sciences

**Table 5. Summary of Kentucky Alfalfa Grazing trials, 1994-2006 (stand persistence shown as a percent of the grazing tolerant Alfagraze).**

Variety/Proprietor		Variety Characteristics <sup>1</sup>						1994 <sup>3,4</sup> 3yr <sup>6</sup>	1996 3yr	1997 4yr	1998 3yr	2000 2yr	2000 3yr	2001 3yr	2004 2yr	Mean <sup>5</sup> (# trials)
		FD	Disease Resistance <sup>2</sup>													
			Bw	Fw	An	PRR	APH									
ABT 205	W-L Research	2	HR	HR	HR	HR	R	94		84						89(2)
ABT 350	W-L Research	3	HR	HR	HR	HR	HR					46				-
ABT 405	W-L Research	4	HR	HR	HR	HR	R	71	129	69		46	100		83(5)	
Alfagraze	Americas Alfalfa	2	MR	R	MR	R	-	100	100	100	100	100	100	100	100(8)	
Amerigraze 401+Z	Americas Alfalfa	4	HR	HR	HR	HR	R		120	53	56	26	85	125	78(6)	
Apollo	Americas Alfalfa	4	R	R	R	R	-	48	75	33	47	17	31	25	39(7)	
Arc (certified)	Public	4	LR	MR	HR	-	-		38						-	
Baralfa 54	Barenbrug USA	-	R	HR	HR	HR	HR				78				-	
Cut-n-Graze	Americas Alfalfa	3	HR	HR	HR	HR	R	68							-	
FK 421	Donley Seed Co.	4	HR	H	H	H	H						100		-	
Feast	Garst Seeds	3	HR	HR	HR	HR	R		146			87	92		108(3)	
Fortress	Syngenta	3	R	R	R	HR	R	40	71						56(2)	
Gold Plus	PGI Alfalfa	4	HR	HR	HR	HR	R				81				-	
Grazeking	FFR/Southern States	5	MR	HR	HR	R	S		91	41			50		61(3)	
Haygrazer	Great Plains Research	4	HR	HR	R	R	MR		75	39			38		51(3)	
Legacy	Green Seed	4	R	R	R	R	R	32							-	
Magnagraze	Dairyland Seed Co.	3	HR	HR	R	HR	-	56							-	
Pasture Plus	MBS	3	HR	HR	R	HR	MR	60							-	
Pioneer 98	Pioneer	3	HR	R	HR	R	-				56				-	
ProGro	MBS Inc.	4	HR	HR	R	HR	MR				81				-	
Quantum	ABI Alfalfa	2	HR	HR	HR	HR	R	71							-	
Rushmore	Syngenta	4	HR	HR	HR	HR	HR	32							-	
Saranac AR (cert.)	Public	4	MR	R	HR	LR	-		77				100		89(2)	
Spredor 3	Syngenta	1	HR	HR	R	MR	S	71	123		75				90(3)	
Stampede	Allied Seed	3	HR	R	R	HR	R		73						-	
Wintergreen	ABI Alfalfa	3	HR	HR	HR	HR	R	95		57	72				75(3)	
WL 326GZ	W-L Research	4	HR	HR	HR	HR	HR		118		88				103(2)	
115 Brand	Monsanto	3	HR	HR	R	HR	R					56	85		71(2)	
5373	Pioneer	4	HR	HR	HRT	MR	LR	21							-	
5432	Pioneer	4	HR	HR	-	MR	-							56	-	

<sup>1</sup> Variety characteristics: FD=fall dormancy, Bw=bacterial wilt, Fw=fusarium wilt, An=anthracnose, PRR=phytophthora root rot, APH-aphanomyces root rot. Information provided by seed companies.

<sup>2</sup> Disease resistance: S=susceptible, LR=low resistance, MR=moderate resistance, R=resistance, HR=high resistance.

<sup>3</sup> Year trial was established.

<sup>4</sup> Use this summary table as a guide in making variety decisions, but refer to specific yearly reports to determine statistical differences in stand persistence between varieties. To find actual persistence ratings, look in the yearly report for the final year of each specific test. For example, the Lexington trial planted in 1996 was grazed for three years so final persistence report would be "1999 Alfalfa Grazing Tolerance Report" archived in the Kentucky Forage Web site at <[www.uky.edu/Ag/Forage](http://www.uky.edu/Ag/Forage)>.

<sup>5</sup> Mean only presented when respective variety was included in two or more trials.

<sup>6</sup> Number of years of data.



Mention or display of a trademark, proprietary product, or firm in text or figures does not constitute an endorsement and does not imply approval to the exclusion of other suitable products or firms.